

# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US05/006645

International filing date: 01 March 2005 (01.03.2005)

Document type: Certified copy of priority document

Document details: Country/Office: US  
Number: 60/549,019  
Filing date: 01 March 2004 (01.03.2004)

Date of receipt at the International Bureau: 31 March 2005 (31.03.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

1300271

# THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

*March 23, 2005*

**THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.**

**APPLICATION NUMBER: 60/549,019**

**FILING DATE: *March 01, 2004***

**RELATED PCT APPLICATION NUMBER: *PCT/US05/06645***



Certified by

Under Secretary of Commerce  
for Intellectual Property  
and Director of the United States  
Patent and Trademark Office

# PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(b)(2)

CERTIFICATE OF MAILING BY EXPRESS MAIL UNDER 37 C.F.R. §1.10

NUMBER OF EXPRESS MAIL MAILING LABEL **EV325929687US** DATE OF DEPOSIT WITH POSTAL SERVICE **March 1, 2004**

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee service under 37 C.F.R. §1.53(b)(2) on the date indicated above, and is addressed to Mail Stop Provisional Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

Name of Person Mailing Paper or Fee  
**Garen Burgoon**

Signature of Person Mailing Paper or Fee  
*Garen Burgoon*

Docket No. **1282-P03383US00**

Type a plus sign (+) inside this box +

INVENTOR(S)/APPLICANT(S)			
LAST NAME	FIRST NAME	MIDDLE INITIAL	RESIDENCE (CITY AND EITHER STATE OR
CONDON	Stephen	M.	Glenmoore, PA
JACKSON	Randy	William	Glenmoore, PA
LAPORTE	Matthew	G.	Honey Brook, PA
BURNS	Christopher	J.	Malvern, PA
HERBERTZ	Torsten		Honey Brook, PA
GABOURY	Janet	A.	Blue Bell, PA
<b>TITLE OF THE INVENTION (200 characters max)</b> <b>PYRANOINDOLE DERIVATIVES AND THE USE THEREOF FOR THE TREATMENT OF HEPATITIS C VIRUS INFECTION OR DISEASE</b>			
<b>CORRESPONDENCE ADDRESS</b> CUSTOMER NUMBER 000110 Telephone (215) 563-4100 Facsimile (215) 563-4044 Direct Inquiries to: Patrick J. Hagan, Esq.			
STATE <b>PA</b>	ZIP CODE <b>19103</b>	COUNTRY <b>US</b>	
<b>ENCLOSED APPLICATION PARTS (check all that apply)</b> <input checked="" type="checkbox"/> Specification      Number of Pages <u>100</u> <input checked="" type="checkbox"/> Claims      Number of Claims <u>21</u> <input type="checkbox"/> Drawings(s)      Number of Sheets <u>0</u> Other (specify) <u>Abstract - 1 page</u>			
<b>METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT (check one)</b> <input checked="" type="checkbox"/> A check or money order is enclosed to cover the filing fees. <input checked="" type="checkbox"/> In the event a fee is required and not enclosed, or the check is improper, or the fee calculation is in error, the Commissioner is hereby authorized to charge any payment to the account of the undersigned attorneys, Deposit Account No. 04-1406. A duplicate copy of this sheet is enclosed.			
		<b>PROVISIONAL FILING FEE AMOUNT(S) \$160.00</b>	
<input type="checkbox"/> Applicant claims large entity status			

The Invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

☒ No

☐ Yes, the name of the U.S. Government agency contract number is: \_\_\_\_\_

Respectfully submitted  
**DANN, DORFMAN, HERRELL AND SKILLMAN**  
 A Professional Corporation

By

*Patrick J. Hagan*  
 Patrick J. Hagan, Esq.

PTO Reg. No. 27, 643

☐ Additional inventors are being named on separately numbered sheets attached hereto



60/549019



18351 U.S. PTO

## FEE TRANSMITTAL

Complete if known

Application Number: **Not yet assigned**Filing Date: **Concurrently herewith**First Named Inventor: **Stephen M. Condon et al.**Title: **PYRANOINDOLE DERIVATIVES AND THE USE THEREOF FOR THE TREATMENT OF HEPATITIS C VIRUS INFECTION OR DISEASE**TOT. AMT. OF PAYMENT: (1) + (2) + (3) = \$ 160.00Our File No.: **1282-P03383US0**

METHOD OF PAYMENT (check one)	FEE CALCULATION (continued)																																																						
<p>1. The Commissioner is hereby authorized to:</p> <p><input type="checkbox"/> Charge indicated fees</p> <p><input checked="" type="checkbox"/> Charge additional fees</p> <p><input checked="" type="checkbox"/> Credit overpayments</p> <p>to the account of DANN, DORFMAN, HERRELL AND SKILLMAN</p> <p>Deposit Account Number <u>04-1406</u></p> <p>2. Payment enclosed:</p> <p>Check in the amount of \$ <u>160.00</u></p>	<p>3. ADDITIONAL FEES</p> <table style="width: 100%;"> <thead> <tr> <th style="text-align: left;">Fee Description</th> <th style="text-align: right;">Fee Paid</th> </tr> </thead> <tbody> <tr><td>Surcharge-late filing fee or oath</td><td style="text-align: right;">_____</td></tr> <tr><td>Surcharge - late provisional filing fee or cover sheet</td><td style="text-align: right;">_____</td></tr> <tr><td>Non-English specification</td><td style="text-align: right;">_____</td></tr> <tr><td>For filing a request for reexamination</td><td style="text-align: right;">_____</td></tr> <tr><td>Requesting publication of SIR prior to Examiner action</td><td style="text-align: right;">_____</td></tr> <tr><td>Requesting publication of SIR after Examiner action</td><td style="text-align: right;">_____</td></tr> <tr><td>Extension for response within first month</td><td style="text-align: right;">_____</td></tr> <tr><td>Extension for response within second month</td><td style="text-align: right;">_____</td></tr> <tr><td>Extension for response within third month</td><td style="text-align: right;">_____</td></tr> <tr><td>Extension for response within fourth month</td><td style="text-align: right;">_____</td></tr> <tr><td>Notice of Appeal</td><td style="text-align: right;">_____</td></tr> <tr><td>Filing a brief in support of an appeal</td><td style="text-align: right;">_____</td></tr> <tr><td>Request for oral hearing</td><td style="text-align: right;">_____</td></tr> <tr><td>Petition to institute a public use proceeding</td><td style="text-align: right;">_____</td></tr> <tr><td>Petition to revive unavoidably abandoned application</td><td style="text-align: right;">_____</td></tr> <tr><td>Petition to revive unintentionally abandoned application</td><td style="text-align: right;">_____</td></tr> <tr><td>Issue fee</td><td style="text-align: right;">_____</td></tr> <tr><td>Advance Order (10 copies)</td><td style="text-align: right;">_____</td></tr> <tr><td>Publication Fee</td><td style="text-align: right;">_____</td></tr> <tr><td>Petitions to the Commissioner</td><td style="text-align: right;">_____</td></tr> <tr><td>Petitions related to provisional applications</td><td style="text-align: right;">_____</td></tr> <tr><td>Submission of Information Disclosure Stmt.</td><td style="text-align: right;">_____</td></tr> <tr><td>Recording each patent assignment per property (times number of properties)</td><td style="text-align: right;">_____</td></tr> <tr><td>Filing a submission after final rejection (37 CFR 1.129(a))</td><td style="text-align: right;">_____</td></tr> <tr><td>For each additional invention to be examined (37 CFR 1.129(b))</td><td style="text-align: right;">_____</td></tr> <tr><td>Other fee (specify) _____</td><td style="text-align: right;">_____</td></tr> </tbody> </table>	Fee Description	Fee Paid	Surcharge-late filing fee or oath	_____	Surcharge - late provisional filing fee or cover sheet	_____	Non-English specification	_____	For filing a request for reexamination	_____	Requesting publication of SIR prior to Examiner action	_____	Requesting publication of SIR after Examiner action	_____	Extension for response within first month	_____	Extension for response within second month	_____	Extension for response within third month	_____	Extension for response within fourth month	_____	Notice of Appeal	_____	Filing a brief in support of an appeal	_____	Request for oral hearing	_____	Petition to institute a public use proceeding	_____	Petition to revive unavoidably abandoned application	_____	Petition to revive unintentionally abandoned application	_____	Issue fee	_____	Advance Order (10 copies)	_____	Publication Fee	_____	Petitions to the Commissioner	_____	Petitions related to provisional applications	_____	Submission of Information Disclosure Stmt.	_____	Recording each patent assignment per property (times number of properties)	_____	Filing a submission after final rejection (37 CFR 1.129(a))	_____	For each additional invention to be examined (37 CFR 1.129(b))	_____	Other fee (specify) _____	_____
Fee Description	Fee Paid																																																						
Surcharge-late filing fee or oath	_____																																																						
Surcharge - late provisional filing fee or cover sheet	_____																																																						
Non-English specification	_____																																																						
For filing a request for reexamination	_____																																																						
Requesting publication of SIR prior to Examiner action	_____																																																						
Requesting publication of SIR after Examiner action	_____																																																						
Extension for response within first month	_____																																																						
Extension for response within second month	_____																																																						
Extension for response within third month	_____																																																						
Extension for response within fourth month	_____																																																						
Notice of Appeal	_____																																																						
Filing a brief in support of an appeal	_____																																																						
Request for oral hearing	_____																																																						
Petition to institute a public use proceeding	_____																																																						
Petition to revive unavoidably abandoned application	_____																																																						
Petition to revive unintentionally abandoned application	_____																																																						
Issue fee	_____																																																						
Advance Order (10 copies)	_____																																																						
Publication Fee	_____																																																						
Petitions to the Commissioner	_____																																																						
Petitions related to provisional applications	_____																																																						
Submission of Information Disclosure Stmt.	_____																																																						
Recording each patent assignment per property (times number of properties)	_____																																																						
Filing a submission after final rejection (37 CFR 1.129(a))	_____																																																						
For each additional invention to be examined (37 CFR 1.129(b))	_____																																																						
Other fee (specify) _____	_____																																																						
<p style="text-align: center;"><b>FEE CALCULATION</b></p> <p>1. FILING FEE</p> <table style="width: 100%;"> <thead> <tr> <th style="text-align: left;">Fee Description</th> <th style="text-align: right;">Fee Paid</th> </tr> </thead> <tbody> <tr><td>Utility filing fee</td><td style="text-align: right;">_____</td></tr> <tr><td>Design filing fee</td><td style="text-align: right;">_____</td></tr> <tr><td>Plant filing fee</td><td style="text-align: right;">_____</td></tr> <tr><td>Reissue filing fee</td><td style="text-align: right;">_____</td></tr> <tr><td>Provisional filing fee</td><td style="text-align: right;">160.00</td></tr> <tr> <td style="text-align: right;"><b>SUBTOTAL (1)</b></td> <td style="text-align: right;"><b>\$ 160.00</b></td> </tr> </tbody> </table> <p>2. CLAIMS</p> <table style="width: 100%;"> <thead> <tr> <th></th> <th style="text-align: center;">Extra</th> <th style="text-align: center;">Fee</th> <th style="text-align: center;">Fee Paid</th> </tr> </thead> <tbody> <tr> <td>Total Claims Presented _____</td> <td style="text-align: center;">20</td> <td style="text-align: center;">= 0</td> <td style="text-align: center;">x 9.00/18.00 = _____</td> </tr> <tr> <td colspan="4" style="text-align: center;">(a)</td> </tr> <tr> <td>Independent Claims Presented _____</td> <td style="text-align: center;">3</td> <td style="text-align: center;">= 0</td> <td style="text-align: center;">x 42.00/84.00 = _____</td> </tr> <tr> <td colspan="4" style="text-align: center;">(b)</td> </tr> <tr> <td>Multiple Dependent Claim (first presentation) _____</td> <td></td> <td style="text-align: center;">= 0</td> <td></td> </tr> <tr> <td colspan="4">           (a) Enter 20 or number previously paid for            (b) Enter 3 or number previously paid for         </td> </tr> <tr> <td style="text-align: right;"><b>SUBTOTAL (2)</b></td> <td></td> <td></td> <td style="text-align: right;"><b>\$ 0.00</b></td> </tr> </tbody> </table>	Fee Description	Fee Paid	Utility filing fee	_____	Design filing fee	_____	Plant filing fee	_____	Reissue filing fee	_____	Provisional filing fee	160.00	<b>SUBTOTAL (1)</b>	<b>\$ 160.00</b>		Extra	Fee	Fee Paid	Total Claims Presented _____	20	= 0	x 9.00/18.00 = _____	(a)				Independent Claims Presented _____	3	= 0	x 42.00/84.00 = _____	(b)				Multiple Dependent Claim (first presentation) _____		= 0		(a) Enter 20 or number previously paid for (b) Enter 3 or number previously paid for				<b>SUBTOTAL (2)</b>			<b>\$ 0.00</b>	<p style="text-align: right;"><b>SUBTOTAL (3) \$ 160.00</b></p>								
Fee Description	Fee Paid																																																						
Utility filing fee	_____																																																						
Design filing fee	_____																																																						
Plant filing fee	_____																																																						
Reissue filing fee	_____																																																						
Provisional filing fee	160.00																																																						
<b>SUBTOTAL (1)</b>	<b>\$ 160.00</b>																																																						
	Extra	Fee	Fee Paid																																																				
Total Claims Presented _____	20	= 0	x 9.00/18.00 = _____																																																				
(a)																																																							
Independent Claims Presented _____	3	= 0	x 42.00/84.00 = _____																																																				
(b)																																																							
Multiple Dependent Claim (first presentation) _____		= 0																																																					
(a) Enter 20 or number previously paid for (b) Enter 3 or number previously paid for																																																							
<b>SUBTOTAL (2)</b>			<b>\$ 0.00</b>																																																				

Submitted By:

Typed or

Printed Name Patrick J. Hagan Reg. Number 27,643

Signature

Patrick J. HaganDate March 1, 2004

Deposit Account User ID

04-1406

**PYRANOINDOLE DERIVATIVES AND THE USE THEREOF  
FOR THE TREATMENT OF HEPATITIS C VIRUS INFECTION OR DISEASE**

**BACKGROUND OF THE INVENTION**

Hepatitis C is a common viral infection that can lead to chronic Hepatitis, cirrhosis, liver failure, and hepatocellular carcinoma. Infection with the Hepatitis C virus (HCV) leads to chronic Hepatitis in at least 85% of cases, is the leading reason for liver transplantation, and is responsible for at least 10,000 deaths annually in the United States (Hepatology, 1997, 26 (Suppl. 1), 2S-10S).

The Hepatitis C virus is a member of the *Flaviviridae* family, and the genome of HCV is a single-stranded linear RNA of positive sense (Hepatology, 1997, 26 (Suppl. 1), 11S-14S). HCV displays extensive genetic heterogeneity; at least 6 genotypes and more than 50 subtypes have been identified.

There is no effective vaccine to prevent HCV infection. The only therapy currently available is treatment with interferon- $\alpha$  (INF- $\alpha$  or combination therapy of INF- $\alpha$  with the nucleoside analog ribavirin (Antiviral Chemistry and Chemotherapy, 1997, 8, 281-301). However, only about 40% of treated patients develop a sustained response, so there is a need for more effective anti-HCV therapeutic agents.

The HCV genome contains a number of non-structural proteins: NS2, NS3, NS4A, NS4B, NS5A, and NS5B (J. General Virology, 2000, 81, 1631-1648). NS5B is an RNA-dependent RNA polymerase which is essential for viral replication, and therefore, the inhibition of NS5B is a suitable target for the development of therapeutic agents.

In the following US patents, pyranoindole derivatives are disclosed and the compounds are stated to have antidepressant and antiulcer activity: 3,880,853 (4/29/75), 4,118,394 (10/3/78). In US patent 4,179,503 (12/18/78) pyranoindoles are disclosed and stated to have diuretic activity. In the following US patents, pyranoindole derivatives are disclosed and the compounds are stated to have antiinflammatory, analgesic, antibacterial, and antifungal activity: 3,843,681

(10/22/74), 3,939,178 (2/17/76), 3,974,179 (8/10/76), 4,070,371 (1/24/78), 4,076,831 (2/28/78).

In the following US patents, pyranoindole derivatives are disclosed and the compounds are stated to have antiinflammatory and analgesic activity: 4,670,462 (6/2/87), 4,686,213 (8/11/87), 4,785,015 (11/15/88), 4,810,699 (3/7/89), 4,822,781 (4/18/89), 4,960,902 (10/2/90). In US patent 5,776,967 (7/7/98) and US patent 5,830,911 (11/3/98), pyranoindole derivatives are disclosed and the compounds are said to inhibit cyclooxygenase-2 and be useful for treating arthritic disorders, colorectal cancer, and Alzheimer's disease.


Also, in the following US patents, processes for preparing pyranoindole derivatives are disclosed: 4,012,417 (3/15/77), 4,036,842 (7/19/77), 4,585,877 (4/29/86), 4,822,893 (4/18/89). Processes for the resolution of racemic pyranoindole derivatives are disclosed in the following US patents: 4,501,899 (2/26/85), 4,515,961 (5/7/85), 4,520,203 (5/28/85), 4,544,757 (10/1/85).

Provisional Application No. 60/382,148 (filed on May 21, 2002) discloses methods of using pyranoindole compounds to treat infection with Hepatitis C. Provisional Application No. 60/382,154 (filed on May 21, 2002) discloses pyranoindole compositions useful for the treatment of Hepatitis C Infection or Disease.

## BRIEF SUMMARY OF THE INVENTION

This invention relates to pyranoindole derivatives, processes for their preparation and pharmaceutical compositions containing them, and to their use in the treatment of Hepatitis C viral infection.

In accordance with this invention there is provided a group of compounds represented by Formula I:



I

wherein:

R<sub>1</sub> is hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, or an arylalkyl or an alkylaryl of 7 to 12 carbon atoms;

R<sub>2</sub> is hydrogen, a straight chain alkyl of 1 to 12 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, an alkoxyalkyl of 2 to 12 carbon atoms, an arylalkyl or alkylaryl of 7 to 12 carbon atoms, a cyanoalkyl of 1 to 8 carbon atoms, an alkylthioalkyl of 2 to 16 carbon atoms, a cycloalkyl-alkyl of 4 to 24 carbon atoms, a substituted or unsubstituted aryl, or a heteroaryl;

R<sub>3</sub> - R<sub>6</sub> are independently hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, a substituted or unsubstituted aryl, furanylmethyl, arylalkyl or alkylaryl of 7 to 12 carbon atoms, alkynyl of 2 to 7 carbon atoms, or R<sub>5</sub> and R<sub>6</sub> together with the ring carbon atom to which they are attached form a carbonyl group;

R<sub>7</sub>- R<sub>8</sub> and R<sub>10</sub> are independently hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, a substituted or unsubstituted aryl, a substituted or

unsubstituted heteroaryl, furanylmethyl, arylalkyl or alkylaryl of 7 to 12 carbon atoms, alkynyl of 2 to 7 carbon atoms, phenylalkynyl, alkoxy of 1 to 8 carbon atoms, arylalkoxy of 7 to 12 carbon atoms, alkylthio of 1 to 8 carbon atoms, trifluoromethoxy, trifluoroethoxy, trifluoromethylthio, trifluoroethylthio, acyl of 1 to 6 carbon atoms, COOH, COO-alkyl, CONR<sub>11</sub>R<sub>12</sub>, F, Cl, Br, I, CN, CF<sub>3</sub>, NO<sub>2</sub>, alkylsulfinyl of 1 to 8 carbon atoms, alkylsulfonyl of 1 to 6 carbon atoms, pyrrolidinyl, or thiazolidinyl;

R<sub>9</sub> is hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, a cycloalkyl of 3 to 12 carbon atoms, a cycloalkyl-alkyl of 4 to 24 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, an alkoxyalkyl of 2 to 12 carbon atoms, an alkoxyalkoxyalkyl of 3 to 18 carbon atoms, an arylalkoxyalkyl of 3 to 18 carbon atoms, a cycloalkylalkoxyalkyl of 3 to 18 carbon atoms, an aryloxyalkyl of 3 to 18 carbon atoms, a heteroaryloxyalkyl of 3 to 18 carbon atoms, an arylthioalkyl of 3 to 18 carbon atoms, a heteroarylthioalkyl of 3 to 18 carbon atoms, a hydroxyalkyl of 1 to 12 carbon atoms, an alkoxyiminoalkyl of 2 to 16 carbon atoms, an alkylthioalkyl of 2 to 16 carbon atoms, an alkylsulfonylalkyl group of 2 to 16 carbon atoms, a monoalkylaminoalkyl of 2 to 16 carbon atoms, a dialkylaminoalkyl of 3 to 16 carbon atoms, a substituted dialkylaminoalkyl of 3 to 16 carbon atoms, a substituted or unsubstituted aryl, arylalkyl of 7 to 12 carbon atoms, a substituted or unsubstituted heteroaryl of 7 to 12 carbon atoms, a substituted or unsubstituted heteroarylalkyl, a substituted or unsubstituted heterocyclic group, and a heterocycle-alkyl;

R<sub>11</sub> - R<sub>12</sub> are independently H, straight chain alkyl of 1 to 8 carbon atoms, branched alkyl of 3 to 12 carbon atoms, cycloalkyl of 3 to 12 carbon atoms, a substituted or unsubstituted aryl or heteroaryl;


M is a bond, CH<sub>2</sub>, or CH<sub>2</sub>CH<sub>2</sub>, with the proviso that when M is a bond, then R<sub>9</sub> is other than a hydroxyl, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, or an arylalkyl;



Y is a bond, CH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>, aryl, or R<sub>2</sub> and Y together with the ring carbon atom to which they are attached may additionally form a spirocyclic cycloalkyl ring of 3 to 8 carbon atoms; or

a crystalline form or a pharmaceutically acceptable salt thereof.

Preferred compounds of the invention include the compounds of Formula I having the formula:



wherein R<sub>1</sub> through R<sub>10</sub> are as defined above.

A preferred embodiment of the invention include the compounds of Formula I wherein:

R<sub>1</sub>-R<sub>8</sub>, R<sub>10</sub>, and Y are as defined above for compounds of Formula I;

R<sub>9</sub> is hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, a cycloalkyl-alkyl of 4 to 24 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, an alkoxyalkyl of 2 to 12 carbon atoms, an alkoxyalkoxyalkyl of 3 to 18 carbon atoms, an arylalkoxyalkyl of 3 to 18 carbon atoms, a cycloalkylalkoxyalkyl of 3 to 18 carbon atoms, an aryloxyalkyl of 3 to 18 carbon atoms, a heteroaryloxyalkyl of 3 to 18 carbon atoms, an arylthioalkyl of 3 to 18 carbon atoms, a heteroarylthioalkyl of 3 to 18 carbon atoms, a hydroxyalkyl of 1 to 12 carbon atoms, an alkoxyiminoalkyl of 2 to 16 carbon atoms, an alkylthioalkyl of 2 to 16 carbon atoms, an alkylsulfonylalkyl group of 2 to 16 carbon atoms, a monoalkylaminoalkyl of 2 to 16 carbon

atoms, a dialkylaminoalkyl of 3 to 16 carbon atoms, a substituted dialkylaminoalkyl of 3 to 16 carbon atoms, a substituted or unsubstituted aryl, arylalkyl of 7 to 12 carbon atoms, a substituted or unsubstituted heteroaryl of 7 to 12 carbon atoms, a substituted or unsubstituted heteroarylalkyl, a substituted or unsubstituted heterocyclic group, and a heterocycle-alkyl;

and M is  $\text{CH}_2$  or  $\text{CH}_2\text{CH}_2$ ; or a crystalline form or a pharmaceutically acceptable salt thereof.

Another preferred embodiment include the compounds of Formula I wherein:

$\text{R}_1$ - $\text{R}_8$ ,  $\text{R}_{10}$ , and Y are as defined above for compounds of Formula 1;

$\text{R}_9$  is a cycloalkyl of 3 to 12 carbon atoms, a cycloalkyl-alkyl of 4 to 24 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, an alkoxyalkyl of 2 to 12 carbon atoms, an alkoxyalkoxyalkyl of 3 to 18 carbon atoms, an arylalkoxyalkyl of 3 to 18 carbon atoms, a cycloalkylalkoxyalkyl of 3 to 18 carbon atoms, an aryloxyalkyl of 3 to 18 carbon atoms, a heteroaryloxyalkyl of 3 to 18 carbon atoms, an arylthioalkyl of 3 to 18 carbon atoms, a heteroarylthioalkyl of 3 to 18 carbon atoms, a hydroxyalkyl of 1 to 12 carbon atoms, an alkoxyiminoalkyl of 2 to 16 carbon atoms, an alkylthioalkyl of 2 to 16 carbon atoms, an alkylsulfonylalkyl group of 2 to 16 carbon atoms, a monoalkylaminoalkyl of 2 to 16 carbon atoms, a dialkylaminoalkyl of 3 to 16 carbon atoms, a substituted dialkylaminoalkyl of 3 to 16 carbon atoms, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl of 7 to 12 carbon atoms, a substituted or unsubstituted heteroarylalkyl, a substituted or unsubstituted heterocyclic group, and a heterocycle-alkyl;

and M is a bond; or a crystalline form or a pharmaceutically acceptable salt thereof.

For purposes of this invention the term "alkyl" includes both straight and branched alkyl moieties, preferably of 1 to 8 carbon atoms. The term "alkenyl" refers to a radical aliphatic hydrocarbon containing at least one double bond and includes both straight and branched alkenyl

moieties of 2 to 7 carbon atoms. Such alkenyl moieties may exist in the E or Z configurations; the compounds of this invention include both configurations. The term "alkynyl" includes both straight chain and branched moieties containing 2 to 7 carbon atoms having at least one triple bond.

The term "cycloalkyl" refers to alicyclic hydrocarbon groups having 3 to 12 carbon atoms and includes but is not limited to: cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, norbornyl, and adamantyl.

For purposes of this invention the term "aryl" is defined as an aromatic hydrocarbon moiety and may be substituted or unsubstituted. An aryl may be selected from but not limited to, the group: phenyl,  $\alpha$ -naphthyl,  $\beta$ -naphthyl, biphenyl, anthryl, tetrahydronaphthyl, phenanthryl, fluorenyl, indanyl, biphenylenyl, acenaphthenyl, acenaphthylenyl, or phenanthrenyl groups. In one embodiment the substituted aryl may be optionally mono-, di-, tri- or tetra-substituted with substituents independently selected from, but not limited to, the group consisting of alkyl, acyl, alkoxycarbonyl, alkoxy, alkoxyalkyl, alkoxyalkoxy, cyano, halogen, hydroxy, nitro, trifluoromethyl, trifluoromethoxy, trifluoropropyl, amino, alkylamino, dialkylamino, amido, dialkylaminoalkyl, alkoxyiminoalkyl, hydroxyalkyl, cycloalkyl, alkylthio,  $-\text{SO}_3\text{H}$ , alkylsulfonyl,  $-\text{SO}_2\text{NH}_2$ ,  $-\text{SO}_2\text{NHalkyl}$ ,  $-\text{SO}_2\text{N(alkyl)}_2$ , alkylsulfonamido, alkenylsulfonamido, alkynylsulfonamido,  $-\text{CO}_2\text{H}$ ,  $-\text{CO}_2\text{NH}_2$ ,  $-\text{CO}_2\text{NHalkyl}$ , and  $-\text{CO}_2\text{N(alkyl)}_2$ . Preferred substituents for aryl include: alkyl, alkoxy, cycloalkyl, acyl, halogen, hydroxyalkyl, amino, alkylamino, dialkylamino, amido, alkylsulfonamido, alkylsulfonyl,  $-\text{CO}_2\text{H}$ ,  $-\text{CO}_2\text{NH}_2$ , trifluoromethyl, trifluoromethoxy, arylalkyl, and alkylaryl.

For purposes of this invention the term "heteroaryl" is defined as an aromatic heterocyclic ring system (monocyclic or bicyclic), which may be substituted or unsubstituted, where the heteroaryl moieties are five or six membered rings containing 1 to 4 heteroatoms selected from the group consisting of S, N, and O, and include but is not limited to: (1) a monocyclic aromatic heterocycle such as furan, thiophene, indole, azaindole, oxazole, thiazole, isoxazole, isothiazole, imidazole, N-methylimidazole, pyridine, pyrimidine, pyrazine, pyrrole, N-methylpyrrole, pyrazole, N-methylpyrazole, 1,3,4-oxadiazole, 1,2,4-oxadiazole, 1,2,4-triazole, 1-methyl-1,2,4-triazole, 1,3,4-thiadiazole, 1,2,4-thiadiazole, 1H-tetrazole, and 1-methyltetrazole; (2) a bicyclic

aromatic heterocycle where a phenyl, pyridine, pyrimidine or pyridazine ring is: (i) fused to a 6-membered aromatic (unsaturated) heterocyclic ring having one nitrogen atom; (ii) fused to a 5 or 6-membered aromatic (unsaturated) heterocyclic ring having two nitrogen atoms; (iii) fused to a 5-membered aromatic (unsaturated) heterocyclic ring having one nitrogen atom together with either one oxygen or one sulfur atom; or (iv) fused to a 5-membered aromatic (unsaturated) heterocyclic ring having one heteroatom selected from O, N or S. Bicyclic aromatic heterocycles include, but are not limited to, benzoxazole, benzothiazole, benzofuran, benzisoxazole, benzimidazole, N-methylbenzimidazole, azabenzimidazole, indazole, quinazoline, quinoline, and pyrrolidine. In one embodiment the substituted heteroaryl may be optionally mono-, di-, tri- or tetra-substituted with substituents independently selected from, but not limited to, the group consisting of alkyl, acyl, alkoxycarbonyl, alkoxy, alkoxyalkyl, alkoxyalkoxy, cyano, halogen, hydroxy, nitro, trifluoromethyl, trifluoromethoxy, trifluoropropyl, amino, alkylamino, dialkylamino, amido, dialkylaminoalkyl, alkoxyiminoalkyl, hydroxyalkyl, cycloalkyl, alkylthio, -SO<sub>3</sub>H, alkylsulfonyl, -SO<sub>2</sub>NH<sub>2</sub>, -SO<sub>2</sub>NHalkyl, -SO<sub>2</sub>N(alkyl)<sub>2</sub>, alkylsulfonamido, alkenylsulfonamido, alkynylsulfonamido, -CO<sub>2</sub>H, -CO<sub>2</sub>NH<sub>2</sub>, -CO<sub>2</sub>NHalkyl, and -CO<sub>2</sub>N(alkyl)<sub>2</sub>. Preferred substituents for heteroaryl include: alkyl, alkoxy, cycloalkyl, acyl, halogen, hydroxyalkyl, amino, alkylamino, dialkylamino, amido, alkylsulfonamido, alkylsulfonyl, -CO<sub>2</sub>H, -CO<sub>2</sub>NH<sub>2</sub>, trifluoromethyl, trifluoromethoxy, arylalkyl, and alkylaryl.

For the purposes of this invention the term “alkoxy” is defined as C1-C12-alkyl-O-; the term “aryloxy” is defined as aryl-O-; the term “heteroaryloxy” is defined as heteroaryl-O-; wherein alkyl, aryl, and heteroaryl are as defined above.

For purposes of this invention the term “arylalkyl” is defined as aryl-C1-C6-alkyl-, and may be substituted or unsubstituted; arylalkyl moieties include benzyl, 1-phenylethyl, 2-phenylethyl, 3-phenylpropyl, 2-phenylpropyl and the like. In one embodiment the substituted arylalkyl may be optionally mono-, di-, tri- or tetra-substituted with substituents independently selected from, but not limited to, the group consisting of alkyl, acyl, alkoxycarbonyl, alkoxy, alkoxyalkyl, alkoxyalkoxy, cyano, halogen, hydroxy, nitro, trifluoromethyl, trifluoromethoxy, trifluoropropyl, amino, alkylamino, dialkylamino, amido, dialkylaminoalkyl, alkoxyiminoalkyl, hydroxyalkyl, cycloalkyl, alkylthio, -SO<sub>3</sub>H, alkylsulfonyl, -SO<sub>2</sub>NH<sub>2</sub>, -SO<sub>2</sub>NHalkyl, -SO<sub>2</sub>N(alkyl)<sub>2</sub>, alkylsulfonamido, -CO<sub>2</sub>H, -CO<sub>2</sub>NH<sub>2</sub>, -CO<sub>2</sub>NHalkyl, and -CO<sub>2</sub>N(alkyl)<sub>2</sub>.

Preferred substituents for arylalkyl include: alkyl, alkoxy, cycloalkyl, acyl, halogen, hydroxyalkyl, amino, alkylamino, dialkylamino, amido, alkylsulfonamido, alkylsulfonyl,  $-\text{CO}_2\text{H}$ ,  $-\text{CO}_2\text{NH}_2$ , trifluoromethyl, trifluoromethoxy, arylalkyl, and alkylaryl.

For purposes of this invention the term “heteroarylalkyl” is defined as heteroaryl-C1-C6-alkyl-, and may be substituted or unsubstituted. In one embodiment the substituted heteroarylalkyl may be optionally mono-, di-, tri- or tetra-substituted with substituents independently selected from, but not limited to, the group consisting of alkyl, acyl, alkoxycarbonyl, alkoxy, alkoxyalkyl, alkoxyalkoxy, cyano, halogen, hydroxy, nitro, trifluoromethyl, trifluoromethoxy, trifluoropropyl, amino, alkylamino, dialkylamino, amido, dialkylaminoalkyl, alkoxyiminoalkyl, hydroxyalkyl, cycloalkyl, alkylthio,  $-\text{SO}_3\text{H}$ , alkylsulfonyl,  $-\text{SO}_2\text{NH}_2$ ,  $-\text{SO}_2\text{NHalkyl}$ ,  $-\text{SO}_2\text{N(alkyl)}_2$ , alkylsulfonamido, morpholine,  $-\text{CO}_2\text{H}$ ,  $-\text{CO}_2\text{NH}_2$ ,  $-\text{CO}_2\text{NHalkyl}$ , and  $-\text{CO}_2\text{N(alkyl)}_2$ . Preferred substituents for heteroarylalkyl include: alkyl, alkoxy, cycloalkyl, acyl, halogen, hydroxyalkyl, amino, alkylamino, dialkylamino, amido, alkylsulfonamido, alkylsulfonyl,  $-\text{CO}_2\text{H}$ ,  $-\text{CO}_2\text{NH}_2$ , morpholine, trifluoromethyl, trifluoromethoxy, arylalkyl, and alkylaryl.

For purposes of this invention the term “alkylaryl” is defined as C1-C6-alkyl-aryl-.

For purposes of this invention the term “alkylthio” is defined as C1-C6-alkyl-S-.

For purposes of this invention “alkoxyalkyl,” “cycloalkyl-alkyl,” “alkylthioalkyl,” “aryloxyalkyl,” and “heteroaryloxyalkyl” denote an alkyl group as defined above that is further substituted with an alkoxy, cycloalkyl, alkylthio, aryloxy, or heteroaryloxy group as defined above.

For purposes of this invention “arylalkoxy,” “alkoxyalkoxy” and “cycloalkylalkoxy” denote an alkoxy group as defined above that is further substituted with an aryl, alkoxy, alkylthio, cycloalkyl or heteroaryl group as defined above.

For purposes of this invention “arylthioalkyl” and “heteroarylthioalkyl” denote an alkyl group as defined above that is further substituted with an arylthio or heteroarylthio group as defined above.

For purposes of this invention “aryloxyalkylthio” is defined as aryloxy-C1-C8-alkyl-S-; “heteroaryloxyalkylthio” is defined as heteroaryloxy-C1-C8-alkyl-S-; where aryloxy, heteroaryloxy, and alkyl are defined above.

For purposes of this invention “alkoxyimino” is defined as alkoxy-N=C-; alkoxyiminoalkyl denotes an alkyl group as defined above that is further substituted with an alkoxyimino group.

For purposes of this invention “arylalkoxyalkyl,” “alkoxyalkoxyalkyl,” and “cycloalkylalkoxyalkyl” denotes an alkyl group as defined above that is further substituted with an arylalkoxy, alkoxyalkoxy, or cycloalkylalkoxy as defined above.

For purposes of this invention “phenylalkynyl” is an alkynyl group further substituted with a phenyl group.

In the most preferred embodiment of this invention a substituted methyl comprises a methyl substituent further substituted with, for example, a furanyl group. In another embodiment of this invention a furanyl substituent is further substituted with a methyl group.

In a preferred embodiment of this invention trifluoromethoxy is  $\text{CF}_3\text{O}-$ . In another embodiment of this invention trifluoromethylthio is  $\text{CF}_3\text{S}-$ .

In one embodiment of this invention trifluoroethoxy includes but is not limited to  $\text{CF}_3\text{CH}_2\text{O}-$ . In another embodiment of this invention trifluoroethylthio includes but is not limited to  $\text{CF}_3\text{CH}_2\text{S}-$ .

The terms “monoalkylamino” and “dialkylamino” refer to moieties with one or two alkyl groups wherein the alkyl chain is 1 to 8 carbons and the groups may be the same or different. The terms monoalkylaminoalkyl and dialkylaminoalkyl refer to monoalkylamino and dialkylamino moieties with one or two alkyl groups (the same or different) bonded to the nitrogen atom which is attached to an alkyl group of 1 to 8 carbon atoms. The terms “substituted monoalkylaminoalkyl” and “substituted dialkylaminoalkyl” refer to monoalkylaminoalkyl and dialkylaminoalkyl moieties that are further substituted with one or more substituents independently selected from the group consisting of aryl.

“Acyl” is a radical of the formula  $-(C=O)$ -alkyl or  $-(C=O)$ -perfluoroalkyl wherein the alkyl radical or perfluoroalkyl radical is 1 to 7 carbon atoms; preferred examples include but are not limited to, acetyl, propionyl, butyryl, trifluoroacetyl.

“Amido” is a radical of the formula  $-NH(C=O)$ -alkyl or  $-N(alkyl)(C=O)$ -alkyl, wherein the alkyl(s) radical are independently 1 to 7 carbon atoms.

For purposes of this invention the term “alkylsulfinyl” is defined as a  $R'SO$ - radical, where  $R'$  is an alkyl radical of 1 to 8 carbon atoms. Alkylsulfonyl is a  $R'SO_2$ - radical, where  $R'$  is an alkyl radical of 1 to 8 carbon atoms. An alkylsulfonylalkyl is an alkyl group of 1 to 8 carbons that is further substituted with an alkylsulfonyl group. Alkylsulfonamido, alkenylsulfonamido, alkynylsulfonamido are  $R'SO_2NR''$ - radicals, where  $R'$  is an alkyl radical of 1 to 8 carbon atoms, an alkenyl radical of 2 to 8 carbon atoms, or an alkynyl radical of 2 to 8 carbon atoms, respectively and  $R''$  is hydrogen or an alkyl radical of 1 to 8 carbon atoms.

For purposes of this invention the term “heterocyclic group” is defined as a heterocyclic ring system (monocyclic or bicyclic) containing at least one saturated or partially saturated heteroaryl group as is defined above, and which may be substituted or unsubstituted. Heterocyclic groups include, but are not limited to (1) a monocyclic saturated or partially saturated heteroaryl group of five or six members containing 1 to 4 heteroatoms selected from the group consisting of S, N, and O; and (2) a bicyclic heterocycle where a phenyl, pyridine, pyrimidine or pyridazine ring is fused to a 5 or 6-membered saturated or partially saturated heteroaryl group containing 1 to 4 heteroatoms selected from the group consisting of S, N, and O. Examples of heterocyclic groups include, but are not limited to, the moieties: azetidiny, 1,4-dioxane, 1,3-dioxolane, hexahydroazepine, piperazine, piperidine, pyrrolidine, morpholine, thiomorpholine, dihydrobenzimidazole, dihydrobenzofuran, dihydrobenzothiène, dihydrobenzoxazole, dihydrofuran, dihydroimidazole, dihydroindole, dihydroisooxazole, dihydroisothiazole, dihydrooxadiazole, dihydrooxazole, dihydropyrrazine, dihydropyrazole, dihydropyridine, dihydropyrimidine, dihydropyrrole, dihydroquinoline, dihydrotetrazole, dihydrothiadiazole, dihydrothiazole, dihydrothien, dihydrotriazole, dihydroazetidine, dihydro-1,4-dioxane, tetrahydrofuran, tetrahydropyran, tetrahydrothien, tetrahydrothiopyran,

tetrahydroquinoline, and tetrahydroisoquinoline. In one embodiment the substituted heteroaryl may be optionally mono-, di-, tri- or tetra-substituted with substituents independently selected from, but not limited to, the group consisting of carbonyl and alkylsulfonyl.

For purposes of this invention, the term “heterocycle-alkyl” denotes an alkyl group as defined above that is further substituted with a heterocyclic group as defined above.

For purposes of this invention, the term “BB7” denotes an RNA-dependent RNA polymerase hepatitis C virus protein sequence which is derived from HCV replicon. A discussion of BB7 and related technology can be found in Blight, K. *et al.* (2000) *Science* 290:1972-1974. BB7 can be licensed from Apath, LLC (893 North Warson Road, Saint Louis Missouri 63141, USA). BB7 is also referred to as Con1 HCV sequence and discussions of Con1 can be found in the following references: Lohmann, V. *et al.* (1999) *Science* 285:110-113; Pietschmann, T. *et al.* (2001) *J. Virol.* 73:1252-1264; Lohmann, V. *et al.* (2001) *J. Virol.* 75:1437-1449.


The compounds of this invention may contain an asymmetric carbon atom and some of the compounds of this invention may contain one or more asymmetric centers and may thus give rise to stereoisomers, such as enantiomers and diastereomers. The stereoisomers of the instant invention are named according to the Cahn-Ingold-Prelog System. While shown without respect to stereochemistry in Formula I, the present invention includes all the individual possible stereoisomers; as well as the racemic mixtures and other mixtures of R and S stereoisomers (scalemic mixtures which are mixtures of unequal amounts of enantiomers) and pharmaceutically acceptable salts thereof. It should be noted that stereoisomers of the invention having the same relative configuration at a chiral center may nevertheless have different R and S designations depending on the substitution at the indicated chiral center.

For compounds of this invention containing two chiral centers, four possible stereoisomers are possible; these four stereoisomers are classified as two racemic pairs of diastereomers. These compounds of the invention may be present as racemic diastereomers which would be designated following the convention described in the 1997 Chemical Abstracts Index Guide, Appendix IV (Columbus, OH) whereas the first cited chiral atom is designated R\* and the next cited chiral atom is designated R\* if it possesses the same chirality as the first cited



stereocenter or S\* if it possesses opposite chirality to the first cited stereocenter. Alternatively, these compounds of the invention may be present as non-racemic mixtures of two diastereomers owing to the existence of a predefined stereocenter. In these instances, the predefined stereocenter is assigned based on the Cahn-Ingold-Prelog System and the undefined stereocenter is designated R\* to denote a mixture of both R and S stereoisomers at this center. Compounds of this invention which possess two chiral centers but which are present as single stereoisomers are described using the Cahn-Ingold-Prelog System.


Based on the chiral center at the C1 carbon position in Formula I, a preferred embodiment of the instant invention is the compound of Formula I(a) shown below:



I(a)

Isomer A

The configuration at C1 in Formula I(a) for purposes of this invention is also referred to as “Isomer A”, and the opposite configuration at C1 is herein defined as “Isomer B” and has the Formula I(b) shown below:




I(b)

## Isomer B

In one embodiment of this invention the compound of the invention is comprised of a ratio of Isomer A to Isomer B of greater than 1:1. In the most preferred embodiment the compound is comprised of 100% Isomer A. In further embodiments the compound is comprised of a ratio of Isomer A to Isomer B of at least about 9:1. In another embodiment the compound is comprised of a ratio of Isomer A to Isomer B of at least about 8:1. Additionally the compound is comprised of a ratio of Isomer A to Isomer B of at least about 7:1.

Another embodiment of this invention is where R<sub>2</sub> of Formula I is a sec-butyl group. In a preferred embodiment, the chiral carbon of the sec-butyl group has an S to R configuration ratio of 1:1. In further embodiments, the chiral carbon of the sec-butyl group has an S to R configuration ratio selected from the group consisting of at least 7:1, at least 8:1, and at least 9:1. In a most preferred embodiment of the invention, the chiral carbon of the sec-butyl group has 100% S configuration.

A preferred aspect of this invention, includes the compounds of Formula I(c):




I(c)

wherein  $R_2$ ,  $R_9$ , M, and Y are as defined for compounds of Formula 1, above.

In further embodiments of the compound of this invention, the compound may be selected from any of the compounds described, *supra*.

The present invention provides a pharmaceutical composition comprising a compound of Formula I:



wherein:

$R_1$  is hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, or an arylalkyl or an alkylaryl of 7 to 12 carbon atoms;

$R_2$  is hydrogen, a straight chain alkyl of 1 to 12 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, an alkoxyalkyl of 2 to 12 carbon atoms, an arylalkyl or alkylaryl of 7 to 12 carbon atoms, a cyanoalkyl of 1 to 8 carbon atoms, an alkylthioalkyl of 2 to 16 carbon atoms, a cycloalkyl-alkyl of 4 to 24 carbon atoms, a substituted or unsubstituted aryl, or a heteroaryl;

$R_3 - R_6$  are independently hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl

of 2 to 7 carbon atoms, a substituted or unsubstituted aryl, furanylmethyl, arylalkyl or alkylaryl of 7 to 12 carbon atoms, alkynyl of 2 to 7 carbon atoms, or R<sub>5</sub> and R<sub>6</sub> together with the ring carbon atom to which they are attached form a carbonyl group;

R<sub>7</sub>- R<sub>8</sub> and R<sub>10</sub> are independently hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, furanylmethyl, arylalkyl or alkylaryl of 7 to 12 carbon atoms, alkynyl of 2 to 7 carbon atoms, phenylalkynyl, alkoxy of 1 to 8 carbon atoms, arylalkoxy of 7 to 12 carbon atoms, alkylthio of 1 to 8 carbon atoms, trifluoromethoxy, trifluoroethoxy, trifluoromethylthio, trifluoroethylthio, acyl of 1 to 6 carbon atoms, COOH, COO-alkyl, CONR<sub>11</sub>R<sub>12</sub>, F, Cl, Br, I, CN, CF<sub>3</sub>, NO<sub>2</sub>, alkylsulfinyl of 1 to 8 carbon atoms, alkylsulfonyl of 1 to 6 carbon atoms, pyrrolidinyl, or thiazolidinyl;


R<sub>9</sub> is hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, a cycloalkyl of 3 to 12 carbon atoms, a cycloalkyl-alkyl of 4 to 24 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, an alkoxyalkyl of 2 to 12 carbon atoms, an alkoxyalkoxyalkyl of 3 to 18 carbon atoms, an arylalkoxyalkyl, a cycloalkylalkoxyalkyl, hydroxyalkyl of 1 to 12 carbon atoms, an alkoxyiminoalkyl of 2 to 16 carbon atoms, an alkylthioalkyl of 2 to 16 carbon atoms, an alkylsulfonylalkyl group of 2 to 16 carbon atoms, a monoalkylaminoalkyl of 2 to 16 carbon atoms, a dialkylaminoalkyl of 3 to 16 carbon atoms, a substituted dialkylaminoalkyl of 3 to 16 carbon atoms, a substituted or unsubstituted aryl, arylalkyl of 7 to 12 carbon atoms, a substituted or unsubstituted heteroaryl of 7 to 12 carbon atoms, a substituted or unsubstituted heteroarylalkyl, a substituted or unsubstituted heterocyclic group, and a heterocycle-alkyl;

R<sub>11</sub> - R<sub>12</sub> are independently H, straight chain alkyl of 1 to 8 carbon atoms, branched alkyl of 3 to 12 carbon atoms, cycloalkyl of 3 to 12 carbon atoms, a substituted or unsubstituted aryl or heteroaryl;

M is a bond, CH<sub>2</sub>, or CH<sub>2</sub>CH<sub>2</sub>, with the proviso that when M is a bond, then R<sub>9</sub> is other than a hydroxyl, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, or an arylalkyl;

Y is a bond, CH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>, aryl, or R<sub>2</sub> and Y together with the ring carbon atom to which they are attached may additionally form a spirocyclic cycloalkyl ring of 3 to 8 carbon atoms; or a crystalline form or a pharmaceutically acceptable salt thereof; and a pharmaceutically acceptable carrier medium.

The present invention also provides a pharmaceutical composition comprising a compound of Formula I(c):



wherein R<sub>2</sub>, R<sub>9</sub>, M, and Y are as defined for compounds of Formula 1, and a pharmaceutically acceptable carrier medium.

Pharmaceutically acceptable salts of the compounds of Formula I having acidic moieties at R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub>, R<sub>9</sub>, or R<sub>10</sub> may be formed from organic and inorganic bases. For example alkali metal salts: sodium, lithium, or potassium and N- tetraalkylammonium salts such as N-tetrabutylammonium salts. Similarly, when a compound of this invention contains a basic moiety at R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub>, R<sub>9</sub>, or R<sub>10</sub>, salts can be formed from organic and inorganic acids. For example salts can be formed from acetic, propionic, lactic, citric, tartaric, succinic, fumaric, maleic, malonic, mandelic, malic, phthalic, hydrochloric, hydrobromic, phosphoric, nitric,

sulfuric, methanesulfonic, naphthalenesulfonic, benzenesulfonic, toluenesulfonic, camphorsulfonic, and similarly known acceptable acids.

In one embodiment, the present invention provides for a method of inhibiting the Hepatitis C RNA-dependent RNA polymerase NS5B. The method comprises contacting a cell with an amount of a compound effective to decrease or prevent NS5B function. The cell may be a mammalian cell and more specifically a human cell. The cell may also be a bacterial cell such as for example *E coli*. The cell may include but is not limited to, a neuronal cell, an endothelial cell, a glial cell, a microglial cell, a smooth muscle cell, a somatic cell, a bone marrow cell, a liver cell, an intestinal cell, a germ cell, a myocyte, a mononuclear phagocyte, an endothelial cell, a tumor cell, a lymphocyte cell, a mesangial cell, a retinal epithelial cell, a retinal vascular cell, a ganglion cell or a stem cell. The cell may be a normal cell, an activated cell, a neoplastic cell, a diseased cell, or an infected cell.

In another embodiment, the present invention provides a method for the treatment or prevention of Hepatitis C infection in a mammal. The present invention accordingly provides to a mammal, a pharmaceutical composition that comprises a compound of this invention in combination or association with a pharmaceutically acceptable carrier. The compound of this invention may be administered as the sole therapeutic agent or in combination with other therapeutically effective compounds or therapies for the treatment or prevention of Hepatitis C viral infection in a mammal.

The compounds are preferably provided orally or subcutaneously. The compounds may be provided by intralesional, intraperitoneal, intramuscular or intravenous injection; infusion; liposome-mediated delivery; topical, nasal, anal, vaginal, sublingual, urethral, transdermal, intrathecal, ocular or otic delivery. In order to obtain consistency in providing the compound of this invention it is preferred that a compound of the invention is in the form of a unit dose. Suitable unit dose forms include tablets, capsules and powders in sachets or vials. Such unit dose forms may contain from 0.1 to 100 mg of a compound of the invention and preferably from 2 to 50 mg. Still further preferred unit dosage forms contain 5 to 25 mg of a compound of the present invention. The compounds of the present invention can be administered orally at a dose range of about 0.01 to 100 mg/kg or preferably at a dose range of 0.1 to 10 mg/kg. Such

compounds may be administered from 1 to 6 times a day, more usually from 1 to 4 times a day. The effective amount will be known to one of skill in the art; it will also be dependent upon the form of the compound. One of skill in the art could routinely perform empirical activity tests to determine the bioactivity of the compound in bioassays and thus determine the appropriate dosage to administer.

The compounds of the invention may be formulated with conventional excipients, such as a filler, a disintegrating agent, a binder, a lubricant, a flavoring agent, a color additive, or a carrier. The carrier may be for example a diluent, an aerosol, a topical carrier, an aqueous solution, a nonaqueous solution or a solid carrier. The carrier may be a polymer or a toothpaste. A pharmaceutically acceptable carrier medium in this invention encompasses any of the standard pharmaceutically accepted carriers, such as phosphate buffered saline solution, acetate buffered saline solution, water, emulsions such as an oil/water emulsion or a triglyceride emulsion, various types of wetting agents, tablets, coated tablets and capsules.

When provided orally or topically, such compounds would be provided to a subject by delivery in different carriers. Typically, such carriers contain excipients such as starch, milk, sugar, certain types of clay, gelatin, stearic acid, talc, vegetable fats or oils, gums, or glycols. The specific carrier would need to be selected based upon the desired method of delivery, for example, phosphate buffered saline (PBS) could be used for intravenous or systemic delivery and vegetable fats, creams, salves, ointments or gels may be used for topical delivery.

The compounds of the present invention may be delivered together with suitable diluents, preservatives, solubilizers, emulsifiers, adjuvants and/or carriers useful in treatment or prevention of Hepatitis C viral infection. Such compositions are liquids or lyophilized or otherwise dried formulations and include diluents of various buffer content (for example, Tris-HCl, acetate, phosphate), pH and ionic strength, additives such as albumins or gelatin to prevent absorption to surfaces, detergents (for example, TWEEN 20, TWEEN 80, PLURONIC F68, bile acid salts), solubilizing agents (for example, glycerol, polyethylene glycerol), anti-oxidants (for example ascorbic acid, sodium metabisulfate), preservatives (for example, thimerosal, benzyl alcohol, parabens), bulking substances or tonicity modifiers (for example, lactose, mannitol), covalent attachment of polymers such as polyethylene glycol, complexation with metal ions, or

incorporation of the compound into or onto particulate preparations of hydrogels or liposomes, micro-emulsions, micelles, unilamellar or multilamellar vesicles, erythrocyte ghosts, or spheroplasts. Such compositions will influence the physical state, solubility, stability, rate of *in vivo* release, and rate of *in vivo* clearance of the compound or composition. The choice of compositions will depend on the physical and chemical properties of the compound capable of treating or preventing a Hepatitis C viral infection.

The compounds of the present invention may be delivered locally via a capsule that allows a sustained release of the compound over a period of time. Controlled or sustained release compositions include formulation in lipophilic depots (for example, fatty acids, waxes, oils).


The present invention further provides a compound of the invention for use as an active therapeutic substance for preventing Hepatitis C infection. Compounds of Formula I are of particular use for the treatment of infection with Hepatitis C virus.

The present invention further provides a method of treating Hepatitis C infection in humans, which comprises administering to the infected individual an effective amount of a compound or a pharmaceutical composition of the invention.

The present invention further provides controlled-release therapeutic dosage forms for the pharmaceutical composition in which the composition is incorporated into a delivery system. The dosage form controls release of the pharmaceutical composition in such a manner that an effective concentration of the composition in the blood can be maintained over an extended period of time, but also the release of the composition should be such that the concentration in the blood remains relatively constant over the extended period of time to improve therapeutic results and/or minimize side effects. Additionally, a controlled release system would affect minimal peak to trough fluctuations in blood plasma levels of the pharmaceutical composition.

The present invention provides a method of treating or preventing a Hepatitis C viral infection in a mammal comprising providing the mammal with an effective amount of at least one pharmaceutical composition, wherein the at least one pharmaceutical composition includes a compound of a formula:





wherein:

R<sub>1</sub> is hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, or an arylalkyl or an alkylaryl of 7 to 12 carbon atoms;

R<sub>2</sub> is hydrogen, a straight chain alkyl of 1 to 12 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, an alkoxyalkyl of 2 to 12 carbon atoms, an arylalkyl or alkylaryl of 7 to 12 carbon atoms, a cyanoalkyl of 1 to 8 carbon atoms, an alkylthioalkyl of 2 to 16 carbon atoms, a cycloalkyl-alkyl of 4 to 24 carbon atoms, a substituted or unsubstituted aryl, or a heteroaryl;

R<sub>3</sub> - R<sub>6</sub> are independently hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, a substituted or unsubstituted aryl, furanylmethyl, arylalkyl or alkylaryl of 7 to 12 carbon atoms, alkynyl of 2 to 7 carbon atoms, or R<sub>5</sub> and R<sub>6</sub> together with the ring carbon atom to which they are attached form a carbonyl group;

R<sub>7</sub>- R<sub>8</sub> and R<sub>10</sub> are independently hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, furanylmethyl, arylalkyl or alkylaryl of 7 to 12 carbon atoms, alkynyl of 2 to 7 carbon atoms, phenylalkynyl, alkoxy of 1 to 8 carbon atoms, arylalkoxy of 7 to 12 carbon atoms, alkylthio of 1 to 8 carbon atoms, trifluoromethoxy,

trifluoroethoxy, trifluoromethylthio, trifluoroethylthio, acyl of 1 to 6 carbon atoms, COOH, COO-alkyl, CONR<sub>11</sub>R<sub>12</sub>, F, Cl, Br, I, CN, CF<sub>3</sub>, NO<sub>2</sub>, alkylsulfinyl of 1 to 8 carbon atoms, alkylsulfonyl of 1 to 6 carbon atoms, pyrrolidinyl, or thiazolidinyl;

R<sub>9</sub> is hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, a cycloalkyl of 3 to 12 carbon atoms, a cycloalkyl-alkyl of 4 to 24 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, an alkoxyalkyl of 2 to 12 carbon atoms, an alkoxyalkoxyalkyl of 3 to 18 carbon atoms, an arylalkoxyalkyl of 3 to 18 carbon atoms, a cycloalkylalkoxyalkyl of 3 to 18 carbon atoms, an aryloxyalkyl of 3 to 18 carbon atoms, a heteroaryloxyalkyl of 3 to 18 carbon atoms, an arylthioalkyl of 3 to 18 carbon atoms, a heteroarylthioalkyl of 3 to 18 carbon atoms, a hydroxyalkyl of 1 to 12 carbon atoms, an alkoxyiminoalkyl of 2 to 16 carbon atoms, an alkylthioalkyl of 2 to 16 carbon atoms, an alkylsulfonylalkyl group of 2 to 16 carbon atoms, a monoalkylaminoalkyl of 2 to 16 carbon atoms, a dialkylaminoalkyl of 3 to 16 carbon atoms, a substituted dialkylaminoalkyl of 3 to 16 carbon atoms, a substituted or unsubstituted aryl, arylalkyl of 7 to 12 carbon atoms, a substituted or unsubstituted heteroaryl of 7 to 12 carbon atoms, a substituted or unsubstituted heteroarylalkyl, a substituted or unsubstituted heterocyclic group, and a heterocycle-alkyl;


R<sub>11</sub> - R<sub>12</sub> are independently H, straight chain alkyl of 1 to 8 carbon atoms, branched alkyl of 3 to 12 carbon atoms, cycloalkyl of 3 to 12 carbon atoms, a substituted or unsubstituted aryl or heteroaryl;

M is a bond, CH<sub>2</sub>, or CH<sub>2</sub>CH<sub>2</sub>, with the proviso that when M is a bond, then R<sub>9</sub> is other than a hydroxyl, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, or an arylalkyl;

Y is a bond, CH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>, aryl, or R<sub>2</sub> and Y together with the ring carbon atom to which they are attached may additionally form a spirocyclic cycloalkyl ring of 3 to 8 carbon atoms; or

a crystalline form or a pharmaceutically acceptable salt thereof.

The present invention also provides a method of treating or preventing a Hepatitis C viral infection in a mammal comprising providing the mammal with an effective amount of at least one pharmaceutical composition, wherein the at least one pharmaceutical composition includes a compound of Formula 1(c):



I(c)

wherein  $R_2$ ,  $R_9$ , M, and Y are as defined for compounds of Formula 1, above.

The method of the present invention further comprises providing the mammal with an effective amount of at least one biologically active agent.

In an embodiment of the method of the present invention, the at least one biologically active agent is provided prior to the at least one pharmaceutical composition, concurrently with the at least one pharmaceutical composition or after the at least one pharmaceutical composition. In a further embodiment of the method of the present invention, the compound is a crystalline form or a pharmaceutically acceptable salt thereof.

In a further embodiment of the method of the present invention, the at least one biologically active agent is selected from the group consisting of interferon, a pegylated interferon, ribavirin, protease inhibitors, polymerase inhibitors, small interfering RNA

compounds, anti-sense compounds, nucleotide analogs, nucleoside analogs, immunoglobulins, immunomodulators, hepatoprotectants, anti-inflammatory agents, antibiotics, antivirals, and anti-infective compounds. In a further embodiment the at least one biologically active agent is a pegylated interferon. In a yet further embodiment the pegylated interferon is a pegylated interferon-alpha.

The compounds of the present invention or precursors thereof and their isomers and pharmaceutically acceptable salts thereof are also useful in treating and preventing viral infections, in particular hepatitis C infection, and diseases in living hosts when used in combination with each other (i.e. pharmaceutical compositions comprising the compounds are administered concurrently with each or sequentially, in either order). The combination of compounds provided herein may further be provided to a subject in respective pharmaceutical compositions, concurrently with or sequentially to other biologically active agents, including but not limited to the group consisting of interferon, a pegylated interferon, ribavirin, protease inhibitors, polymerase inhibitors, small interfering RNA compounds, anti-sense compounds, nucleotide analogs, nucleoside analogs, immunoglobulins, immunomodulators, hepatoprotectants, anti-inflammatory agents, antibiotics, antivirals, and anti-infective compounds. The present invention further provides combination therapy with one or more pyranoindole derivatives, i.e., at least two pharmaceutical compositions, each comprising a different compound of the present invention, are provided to a subject in need thereof either concurrently with each other or sequentially, and such therapy may further comprise providing concurrently or sequentially other medicinal agents or potentiators, such as acyclovir, famicyclovir, valgancyclovir and related compounds, ribavirin and related compounds, amantadine and related compounds, various interferons such as, for example, interferon-alpha, interferon-beta, interferon-gamma and the like, as well as alternative forms of interferons such as pegylated interferons. Additionally, combinations of, for example ribavirin and interferon, may be administered as an additional combination for a multiple combination therapy with at least one of the compounds of the present invention.

The combination therapy with any of the above-described biologically active agents may also be sequential, that is the treatment with a first pharmaceutical composition comprising a compound of the invention followed by treatment with a second pharmaceutical composition

comprising a second compound of the invention, wherein the second compound is different than the first compound; alternatively, treatment may be with both two or more pharmaceutical compositions, wherein each pharmaceutical composition comprises a different compound of the invention, at the same time. The sequential therapy can be within a reasonable time after the completion of the first therapy with the pharmaceutical composition. Treatment with the respective pharmaceutical compositions, each comprising a different compound of the present invention, at the same time may be provided in the same daily dose or in separate doses. Combination therapy may also be provided wherein a pharmaceutical composition comprising at least one compound of the present invention is administered in a composition further comprising at least one biologically active agent, i.e. in a single dose. The dosages for both concurrent and sequential combination therapy (for combined pharmaceutical compositions comprising at least two compounds of the invention or compositions comprising at least one compound of the invention and at least one biologically active agent), will depend on absorption, distribution, metabolism and excretion rates of the components of the pharmaceutical composition as well as other factors known to one of skill in the art. Dosage values of the pharmaceutical composition will also vary with the severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens and schedules may be adjusted over time according to the individual's need and the professional judgment of the person administering or supervising the administration of the pharmaceutical compositions.

In a further embodiment, the compounds of the invention may be used for the treatment of HCV in humans in combination therapy mode with other inhibitors of the HCV polymerase.

In yet a further embodiment, the compounds of the present invention may be used for the treatment of HCV in humans in combination therapy mode with other inhibitors of the HCV life cycle such as, for example, inhibitors of HCV cell attachment or virus entry, HCV translation, HCV RNA transcription or replication, HCV maturation, assembly or virus release, or inhibitors of HCV enzyme activities such as the HCV nucleotidyl transferase, helicase, protease or polymerase.

It is intended that combination therapies of the pharmaceutical compositions include any chemically compatible combination of a compound of this inventive group with other compounds of the inventive group or other compounds outside of the inventive group, as long as the combination does not eliminate the anti-viral activity of the compound of this inventive group or the anti-viral activity of the pharmaceutical composition itself.

The term "interferon-alpha" as used herein means the family of highly homologous species-specific proteins that inhibit viral replication and cellular proliferation and modulate immune response. Typical suitable interferon-alphas include, but are not limited to, recombinant interferon alpha-2b such as INTRON-A INTERFERON available from Schering Corporation, Kenilworth, NJ, recombinant interferon alpha-2a such as Roferon interferon available from Hofman-La Roche, Nutley, NJ, a recombinant interferon alpha-2C, such as BEROFOR ALPHA 2 INTERFERON available from Boehringer Ingelheim Pharmaceutical, Inc., Ridgefield, Conn., interferon alpha-n1, a purified blend of natural alpha interferons such as SUMIFERON available from Sumitomo, Japan or as Wellferon interferon alpha-n1 (INS) available from Glaxo-Wellcome Ltd., London, Great Britain, or a consensus alpha interferon such as those described in U.S. Patent Nos. 4,897,471 and 4,695,623 (the contents of which are hereby incorporated by reference in their entireties, specifically examples 7, 8 or 9 thereof) and the specific product available from Amgen, Inc., Newbury Park, Calif., or interferon alpha-n3 a mixture of natural interferons made by Interferon Sciences and available from the Purdue Frederick Co., Norwalk, Conn., under the ALFERON trademark. The use of interferon alpha-2a or alpha 2b is preferred. Since interferon alpha 2b, among all interferons, has the broadest approval throughout the world for treating chronic hepatitis C infection, it is most preferred. The manufacture of interferon alpha 2b is described in U.S. Pat. No. 4,503,901.


The term "pegylated interferon" as used herein means polyethylene glycol modified conjugates of interferon, preferably interferon alpha-2a and alpha-2b. The preferred polyethylene-glycol-interferon alpha-2b conjugate is PEG.sub.12000-interferon alpha 2b. The phrase "PEG.sub.12000-IFN alpha" as used herein means conjugates such as are prepared according to the methods of International Application No. WO 95/13090 and containing

urethane linkages between the interferon alpha-2a or alpha-2b amino groups and polyethylene glycol having an average molecular weight of 12000.


## DETAILED DESCRIPTION OF THE INVENTION

The compounds of the present invention can be readily prepared according to the following reaction schemes or modification thereof or otherwise using known chemistry procedures. In the following reaction schemes  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R_6$ ,  $R_7$ ,  $R_8$ ,  $R_9$ , or  $R_{10}$ ,  $R_{11}$ ,  $R_{12}$  and  $Y$  are selected from the groups defined above.


Preferred compounds of the present invention can be synthesized as described in the schemes below (Scheme 1 to 6).




Scheme 1



Scheme 2




Scheme 3




Scheme 4





Scheme 5



Scheme 6

The ability of the compounds of the present invention to inhibit Hepatitis C Polymerase was established by the following experimental procedure:

NS5B from the BK strain (genotype 1b) is expressed in *E coli* as a protein in which the 21 C-terminal amino acids are replaced with a short linker and a hexahistidine tag (GSHHHHHH). The purified protein is mixed with radioactive nucleotides and allowed to replicate a heteropolymeric RNA substrate, primed by an endogenous short hairpin, resulting in an approximately 760 nt product. The radioactive product is captured on a filter and quantitated after removal of the unincorporated nucleotides.

#### Reagents:

10 mM uridine 5'-triphosphate (UTP) (Promega # p116B)

10 mM adenine 5'-triphosphate (ATP) (Promega # p113B)

10 mM cytidine 5'-triphosphate (CTP) (Promega # p114B)  
10 mM guanine 5'-triphosphate (GTP) (Promega # p115B)  
Bovine Serum Albumin (BSA) 10 mg/ml NEB (100X at 10 mg/ml) #007-BSA  
RNasein (Promega #N251X) 40 U/ $\mu$ l  
A-[33P]-GTP (NEN-easytides NEG/606H 3000 Ci/mmol, 370 MBq/ml, 10 mCi/ml)  
Falcon polypropylene 96 well plates (Becton Dickinson # 351190)  
Millipore Multiscreen assay system-96 well-filtration plate #MADE NOB 50  
Optiphase Supermix (Wallac) formulated by Fisher  
Millipore Multiscreen liner for use in microbeta 1450-106 cassette [(Wallac) Perkin Elmer #1450-433]  
1 M (N-[2-hydroxyethyl]piperazine-N'-[2-ethanesulfonic acid]) (HEPES), pH 7.3  
Amersham Pharmacia Biotech (US16924-500 ml)  
1 M  $\text{MgCl}_2$  (SIGMA #M1028)  
Dithiothreitol (DTT) (solid) (SIGMA # D9779)  
RNase free water (GIBCO-BRL #10977-023)  
Dimethyl sulfoxide (Aldrich #27685-5)  
Basilen Blue (Sigma, B5520)  
0.5M ethylenediaminetetraacetic acid (EDTA), pH 8 (GIBCO-BRL #15575-020)  
Dibasic sodium phosphate (7-hydrate) ( $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$ ; Baker#3824-07)  
Phosphoric acid (Baker, #0262.02)

Further reagent preparation:

- 0.5M Na Phosphate buffer. Per liter, weigh 134 gr  $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$ , add water to 900 ml. Adjust pH to 7.0 with phosphoric acid. Top off with water to 1 L.
- Dilute nucleotides 1:1000 to 10  $\mu\text{M}$  (GTP and CTP) or 1:100 to 100  $\mu\text{M}$  (ATP and UTP) into RNase free water.

Procedure:

(1) Compounds 10 $\mu$ l at 10  $\mu$ g/ml in 15 % dimethylsulfoxide (DMSO)

When starting from 100  $\mu$ g/ml compound stock in 1% DMSO:

Dispense 5  $\mu$ l 30 % DMSO per well

Dispense 5  $\mu$ l compound (100  $\mu$ g/ml) per well.

When starting from 50  $\mu$ g/ml compound stock in 15 % DMSO:

Add 10  $\mu$ l compound per well.

(2) Enzyme Mix:

Stock	Final Conc (in 50 $\mu$ l assay volume)	Per 20 $\mu$ l mix (1 reaction)	Per 600 reactions
Diethyl Pyrocarbonate (DEPC) water		17.06 $\mu$ l	10236 $\mu$ l
1 M HEPES, pH 7.5	20 mM	0.5 $\mu$ l	300 $\mu$ l
1 M MgCl <sub>2</sub>	5 mM	0.25 $\mu$ l	150 $\mu$ l
100 mM DTT	1 mM	0.5 $\mu$ l	300 $\mu$ l
100 $\mu$ M UTP	0.5 $\mu$ M	0.25 $\mu$ l	150 $\mu$ l
100 $\mu$ M ATP	1 $\mu$ M	0.5 $\mu$ l	300 $\mu$ l
10 $\mu$ M CTP	0.08 $\mu$ M	0.4 $\mu$ l	240 $\mu$ l
10 $\mu$ M GTP	0.025 $\mu$ M	0.125 $\mu$ l	75 $\mu$ l
BSA, 10 mg/ml	0.05 mg/ml	0.25 $\mu$ l	150 $\mu$ l
HCV RdRp NS5B d21BK (500 $\mu$ g/ml or ~7.5 $\mu$ M)	24 nM	0.16 $\mu$ l	96 $\mu$ l

Total: 20  $\mu$ l

12 ml

Add 20  $\mu$ l enzyme mix into each well of the assay plate. Incubate compound and enzyme at room temperature for 15 minutes

(3) Template mix - prepare ahead

Spin down a tube of RNA (5  $\mu$ g/tube stored in 75% ethanol and 0.3 M sodium acetate) in a microcentrifuge for 20 minutes at 4 °C. One tube is enough for 1 - 1.5 plates. Remove as much ethanol from the tube as possible by inverting the tube. Be gentle, pellet RNA may not adhere to the tube. Vacuum dry the RNA. Resuspend the RNA by adding 1 ml of DEPC water, close the cap of the tube tightly. To dissolve RNA, incubate RNA solution on ice for ~60

minutes and gently vortex. Spin briefly to ensure all RNA solution is down to the bottom of the tube before opening cap. Gently transfer RNA solution into a 5 ml or larger tube. Add another 3 ml of DEPC water (total 4 ml of volume).

Add the following volumes of reagents

Stock	Final concentration	Per 20 µl mix (1 reaction)	Per 600 reactions
RNAse-free water		2.98 µl	1788 µl
HEPES, 1M	20 mM	0.5 µl	300 µl
RNAse Inhibitor (40 U/ µl)	0.4 µ/µl	0.5 µl	300 µl
33P-GTP 3000 Ci/mmol, 10 µCi/ µ l (3.3 µM)	0.025 µM	0.0125 µl	7.5 µl
POF RNA template	3 nM	16 µl	9600 µl

Add 20 µl template mix per reaction (i.e. 20 ng of pOF per reaction or ~3 nM)

(4) Incubate reaction at room temperature (22-25°C) for 2 hours.

(5) Stop reaction by adding 50 µl of 170 mM EDTA.

Final concentration of EDTA is 85 mM.

(6) Prewet filters of Millipore multiscreen assay plate by adding 200 µl of 0.5 M sodium phosphate buffer, pH 7.0 into each well. Let stand at room temperature for 2 - 3 minutes.

(7) Place the multiscreen filter plate onto a Millipore Manifold and turn on vacuum to allow buffer to flow through. Turn off vacuum. Transfer 80 µl of the reaction product into each well of the filter plate. Let stand for 2 - 3 minutes. Turn on vacuum to filter reaction product.

(8) Turn off vacuum. Add 200 µl of 0.5 M sodium phosphate buffer, pH 7.0 into each well to wash filter. Turn on vacuum.

Repeat step (8) three more times.

- (9) Remove polypropylene bottom. Spot dry filter at the bottom with paper towel. Air dry filter plate on a bench for 1 hour. Add 40  $\mu$ l Super Mix scintillant. Seal top of the plate with a tape. Place plate into a Packard carrier or micro-beta carrier.
- (10) Count plate using a Packard Topcount or micro-beta counter. Count (for example using Program 10) for  $^{33}\text{P}$  in Top count or  $^{33}\text{P}$  program in micro-beta.

See, Ferrari et al. 1999. J. Virology 73:1649-1654: "Characterization of soluble Hepatitis C virus RNA-dependent RNA polymerase expressed in E. coli and Takamizawa et al 1991" and J. Virology 65:1105-1113: "Structure and characterization of the Hepatitis C virus genome isolated from human carriers," both references are hereby incorporated by reference.

The compounds of the present invention inhibited Hepatitis C polymerase as summarized in Table 1:

Table 1

<b>Example</b>	<b>BK IC<sub>50</sub> A= &lt;0.5<math>\mu</math>M, B= 0.5 to &lt;5.0 <math>\mu</math>M, C= <math>\geq</math>5 <math>\mu</math>M</b>	<b>BB7 IC<sub>50</sub> A= &lt;0.5<math>\mu</math>M, B= 0.5 to &lt;5.0 <math>\mu</math>M, C= <math>\geq</math>5 <math>\mu</math>M</b>
1	A	A
2	B	A
3	B	A
4	B	A
5	B	A
6	B	A
7	B	B
8	B	A
9	B	B
10	B	B
11	B	B
12	B	B
13	B	B
14	B	B

Example	BK IC <sub>50</sub> A= <0.5 μM, B= 0.5 to <5.0 μM, C= ≥5 μM	BB7 IC <sub>50</sub> A= <0.5 μM, B= 0.5 to <5.0 μM, C= ≥5 μM
15	A	A
16	B	A
17	B	C
18	B	A
19	B	B
20	B	B
21	A	A
22	A	A
23	A	A
24	A	A
25	A	A
26	A	A
27	A	A
28	A	A
29	A	A
30	A	A
31	B	B
32	A	A
33	A	A
34	A	A
35	A	A
36	A	A
37	A	A
38	A	A
39	A	A
40	A	A
41	A	A
42	A	B
43	A	A
44	A	A
45	A	A
46	A	A
47	C	B
48	A	A
49	A	A
50	A	A
51	A	A
52	A	A

Example	<b>BK</b> <b>IC<sub>50</sub></b> A= <0.5 $\mu$ M, B= 0.5 to <5.0 $\mu$ M, C= $\geq$ 5 $\mu$ M	<b>BB7</b> <b>IC<sub>50</sub></b> A= <0.5 $\mu$ M, B= 0.5 to <5.0 $\mu$ M, C= $\geq$ 5 $\mu$ M
53	A	A
54	A	A
55	A	A
56	A	A
57	C	B
58	B	B
59	A	A
60	A	A
61	A	A
62	A	A
63	A	A
64	A	A
65	A	A
66	A	A
67	A	A
68	A	A
69	A	A
70	A	A
71	A	A
72	A	A
73	A	A
74	A	A
75	A	A
76	A	A
77	A	A
78	A	A
79	A	A
80	A	A
81	A	A
82	A	A
83	A	A
84	A	A
85	A	A
86	A	A
87	A	A
88	A	A
89	A	A
90	A	A



Example	BK IC <sub>50</sub> A= <0.5 μM, B= 0.5 to <5.0 μM, C= ≥5 μM	BB7 IC <sub>50</sub> A= <0.5 μM, B= 0.5 to <5.0 μM, C= ≥5 μM
91	A	A
92	A	A
93	A	A
94	A	A
95	A	A
96	A	A
97	A	A
98	B	A
99	A	A
100	A	A
101	A	A
102	A	A
103	A	A
104	A	A
105	A	A
106	A	A
107	A	A
108	A	A
109	A	A
110	A	A
111	A	A
112	A	A
113	A	A
114	A	A
115	A	A
116	A	A
117	A	A
118	A	A
119	A	A
120	A	A
121	A	A
122	A	A
123	A	A
124	A	A
125	A	A
126	A	A
127	A	A
128	A	A

Example	BK IC <sub>50</sub> A= <0.5 μM, B= 0.5 to <5.0 μM, C= ≥5 μM	BB7 IC <sub>50</sub> A= <0.5 μM, B= 0.5 to <5.0 μM, C= ≥5 μM
129	A	A
130	A	A
131	-	-
132	A	A
133	A	A
134	A	A
135	A	A
136	A	A
137	A	A
138	A	A
139	A	A
140	A	A
141	A	A
142	A	A
143	A	A

The ability of the compounds of the present invention to inhibit Hepatitis C virus replicon constitutively expressed in a human liver cell line was established by the following experimental procedure:

Clone A cells (licensed from Apath, LLC) are derived from Huh-7 cells (human hepatoma cell line) and constitutively express the HCV replication proteins with concomitant amplification the HCV replicon (1b) genome. Cells are maintained and passaged in DMEM/10% FCS/1 mg/ml G418 (Geneticin from Gibco #11811-023; other media components as described below in “elisa media”). Care should be taken to maintain cell monolayers at a subconfluent state by 1:3 or 1:4 passages every 3-4 days. The replicon is extremely sensitive to the cellular metabolism/proliferation state and replicon copy number will rapidly decline in confluent monolayers (resting cells). Under ideal conditions each cell has, on average, 1000 copies of the HCV replicon genome.

Reagents:

Elisa media:

Dulbecco's Modified Eagle Media (DMEM) (Gibco #12430-047)

2% Fetal Calf Serum (FCS) (HyClone #SH30070.03)

1X pen/strep (Gibco #15140-122)

1X Non-essential amino acids (NEAA) (Gibco #11140-050)

no G418

Glutaraldehyde (Fisher #02957-4)

TWEEN-20, 10% (Roche #1332465)

TRITON X-100 (Sigma #T-8787)

Superblock in Phosphate Buffered Saline (PBS) (Pierce #37515)

NS5a monoclonal antibody (Virostat #1873)

Goat antimouse-HRP monoclonal antibody (BioRad #172-1011)

3,3',5,5' tetramethylbenzidine (TMB) substrate (Sigma #T-0440)

Compound Dilution/Cell Plating:

Drug Plate Preparation (Mother Plate)

10  $\mu$ l of compounds (in DMSO) are added to column 3 of the mother plate. 5  $\mu$ l of DMSO are added to the remaining columns. Mother plates are set aside until ready for serial dilution to be performed.

Control Drugs

Drug and Cell Addition:

The process for each plate involves:

Prepare cell plates (daughter plates) by adding 52  $\mu$ l of Elisa media to each well.

In Mother plates, serially transfer 50  $\mu$ l/well from column 3 through column 12.

Transfer 8  $\mu$ l from mother plate to daughter plates (all 96 wells).

Place daughter plates in incubator until cells are prepared.

Harvest Clone A cells and plate directly into daughter plates at  $0.7 \times 10^5$  cells/ml, 100  $\mu$ l/well.

All plates are incubated at 37°C in 5% CO<sub>2</sub> for 3 days.

Elisa Assay:

Remove media from 96-well plates (cells should be ca 80% confluent) by flicking into sink.

Add 130  $\mu$ l/well 1X PBS + 0.05% glutaraldehyde.

Incubate 37°C for 1 hour.

Remove by flicking into sink.

Wash 3X with 300  $\mu$ l/well PBS, shaking 5 min each wash. Remove by flicking into sink.

Add 130  $\mu$ l/well PBS + 0.05% TWEEN-20 + 0.1% TRITON X-100.

Incubate 37°C for 10 minutes.

Remove by flicking into sink.

Add 300  $\mu$ l/well Superblock in PBS.

Incubate 37°C for 1 hour.

Remove by flicking into sink.

Wash 3x with 300  $\mu$ l/well PBS, shaking 5 minutes each wash. Remove by flicking into sink.

During last wash, make a 1:100 dilution of NS5a Monoclonal-antibody (Mab) in Superblock + 0.02% TWEEN-20.

After last wash, add 50 µl/well diluted Mab.

Incubate 37°C for 1 hour.

Remove by flicking into sink.

Wash 3X with 300 µl/well PBS + 0.02% TWEEN-20, shaking 5 minutes each wash.

Remove by flicking into sink.

During last wash, make a 1:500 dilution of goat antimouse-HRP Mab in Superblock + 0.02% TWEEN-20.

After last wash, add 50 µl/well diluted Mab.

Incubate 37°C for 1 hour.

Remove by flicking into sink.

Wash 5X with 300 µl/well PBS + 0.02% TWEEN-20, shaking 5 minutes each wash. Remove by flicking into sink.

Wash 3X with 300 µl/well PBS, shaking 5 minutes each wash. Remove by flicking into sink.

After last wash, add 130 µl/well room temperature TMB substrate.

Incubate until blue color develops.

Add 130 µl/well 1N HCl to stop reaction (color turns from blue to yellow).

Read plates with optical density (O.D.) 450 filter.

ANALYSIS OF RESULTS: IC<sub>50</sub> (µM); IC<sub>50</sub> (µg/ml); % Inhibition

REFERENCE COMPOUNDS: Interferon-α<sub>2</sub>; 4-30 U/ml IC<sub>50</sub>

The following non-limiting specific examples are included to illustrate the synthetic procedures used for preparing compounds of the Formula I. In these examples, all chemicals and intermediates are either commercially available or can be prepared by standard procedures found in the literature or are known to those skilled in the art of organic synthesis.

### Example 1

#### **5-Bromo-1-ethoxycarbonylmethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indole-7-carboxylic acid benzyl ester**

A 1000 mL round-bottom flask (RBF) was charged with of 4-bromo-3-(2-hydroxy-ethyl)-7-methyl-1*H*-indole-6-carboxylic acid benzyl ester (12.03 g, 0.031 mol), ethyl butyrylacetate (5.45 mL, 0.034 mol), and 500 mL dichloromethane (DCM). This mixture was cooled to 0°C with stirring. Over approximately 5 minutes boron trifluoride diethyl etherate (BF<sub>3</sub>·Et<sub>2</sub>O) (9.82 mL, 0.078 mol) was added to the stirring mixture. The reaction was allowed to warm to ambient temperature and was stirred for 1 hour. The reaction was then diluted with ethyl acetate (EtOAc), washed saturated NaHCO<sub>3</sub> (2x) and saturated NaCl (1x), dried (MgSO<sub>4</sub>), and concentrated. The resulting yellow oil was purified by flash chromatography on SiO<sub>2</sub> eluting with a 10% EtOAc/hexanes to 20% EtOAc/hexanes gradient. Trituration in hexanes yielded 13.71g (84.0%) of white powder. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 9.81 (br s, 1H), 7.89 (s, 1H), 7.40 (m, 5H), 5.35 (s, 2H), 4.21 (m, 2H), 3.95 (m, 2H), 3.15 (t, J = 4.7 Hz, 2H), 3.00 (m, J = 22.3 Hz, 2H), 2.73 (s, 3H), 2.00 (m, 2H), 1.29 (m, 4H), 0.88 (t, J = 7.6Hz, 3H).

#### **5-Cyano-1-ethoxycarbonylmethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indole-7-carboxylic acid benzyl ester**

A 500 mL RBF was charged with 5-bromo-1-ethoxycarbonylmethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indole-7-carboxylic acid benzyl ester (12.00 g, 23 mmol), 100mL 1-methyl-2-pyrrolidinone (NMP), and CuCN (20.34 g, 0.227 mol). The reaction was heated to 190°C with vigorous stirring for 45 minutes. The reaction was then cooled to ambient temperature and diluted with EtOAc. Water (300mL) was added to the mixture, followed by the addition of a 1:1 mixture of Celite/silica gel. This suspension was stirred for several minutes,

and then filtered through a pad of Celite. The Celite was washed with EtOAc, and the filtrate was washed H<sub>2</sub>O (5x) and saturated NaCl (1x), dried over MgSO<sub>4</sub>, and concentrated. Upon trituration with hexanes, 8.82g (81.8%) of light tan solid was obtained. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 10.11 (br s, 1H), 8.12 (s, 1H), 7.40 (m, 5H), 5.37 (s, 2H), 4.22 (m, 2H), 3.95 (m, 2H), 3.00 (m, 4H), 2.83 (s, 3H), 2.00 (m, 2H), 1.30 (m, 4H), 0.89 (t, J = 7.6 Hz, 3H).

**5-Cyano-1-ethoxycarbonylmethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indole-7-carboxylic acid**

A 500mL Parr flask was charged with 5-cyano-1-ethoxycarbonylmethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indole-7-carboxylic acid benzyl ester (14.24 g, 0.030 mol), 180mL of 1:1 MeOH/EtOAc, and Pd(OH)<sub>2</sub> (20% on C, 3.02 g). The flask was shaken on a Parr shaker at 5psi above ambient pressure for 30 minutes. The reaction mixture was filtered through a pad of Celite, washed with methanol and concentrated. Trituration with hexanes yielded 11.1 g (96.4%) of off-white powder. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 10.21 (br s, 1H), 8.22 (s, 1H), 4.22 (m, 2H), 4.00 (m, 2H), 3.09 (m, 4H), 2.89 (s, 3H), 2.00 (m, 2H), 1.30 (m, 4H), 0.90 (t, J = 7.6 Hz, 3H).

**(5-Cyano-7-hydroxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid ethyl ester**

To a solution of 5-cyano-1-ethoxycarbonylmethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indole-7-carboxylic acid (11.1 g, 29.0 mmol), which can be prepared according to the previous step, in tetrahydrofuran (THF) (150 mL) at 0°C was added BH<sub>3</sub>·THF (1.0M in THF, 72.2 mL, 72.2 mmol). The cooling bath was removed, and the solution was stirred for 1.5 hours. The reaction was quenched with 3% HCl, diluted with EtOAc, and the layers were separated. The organic layer was washed with saturated NaHCO<sub>3</sub>, H<sub>2</sub>O, and brine, dried (MgSO<sub>4</sub>) and concentrated *in vacuo*. Purification via flash chromatography on SiO<sub>2</sub> using 20% ethyl acetate/dichloromethane (EtOAc/DCM) as eluent afforded 8.28 g (77%) of pale green foam. ESI-MS m/z 369 (M-H)<sup>-</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 9.72 (bs, 1H), 7.43 (s, 1H), 4.80 (d, J = 5.5 Hz,

2H), 4.25 (m, 2H), 4.15 (m, 1H), 3.93 (m, 1H), 3.02 (m, 3H), 2.92 (d,  $J = 17.0$  Hz, 1H), 2.57 (s, 3H), 2.00 (m, 2H), 1.30 (m, 5H), 0.88 (t,  $J = 7.6$  Hz, 3H).

## Example 2

### **(5-Cyano-7-methoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid**

### **[5-Cyano-8-methyl-1-propyl-7-(2,2,2-trichloro-acetimidoyloxymethyl)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid ethyl ester**

To a solution of (5-cyano-7-hydroxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid ethyl ester (0.75 g, 2.0 mmol) in DCM (12 mL) was added sodium hydride (9.7 mg, 4.0 mmol). After 10 minutes, trichloroacetonitrile (608  $\mu$ L, 6.0 mmol) was added. The reaction was stirred for 4 hours, at which point additional trichloroacetonitrile (200 mL) was added. After an additional 45 minutes of stirring, sodium hydride (4 mg) was added. The reaction was placed in a freezer at 0°C overnight. After warming to room temperature the next morning, an additional aliquot of sodium hydride (9.7 mg) was added. The reaction was stirred for 1.5 hours, quenched with H<sub>2</sub>O and diluted with EtOAc. The layers were separated, and the organic layer washed with brine, dried (MgSO<sub>4</sub>) and concentrated to afford 1.14 g (109%) of an off-white solid which was used without further purification. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  9.84 (s, 1H), 8.43 (s, 1H), 7.53 (s, 1H), 5.44 (s, 2H), 4.22 (m, 2H), 4.090 (m, 1H), 3.95 (m, 1H), 3.07 (m, 3H), 2.96 (d,  $J = 15.2$  Hz, 1H), 2.61 (s, 3H), 2.06 (m, 1H), 1.95 (m, 1H), 1.42 (m, 1H), 1.12 (m, 4H), 0.91 (t,  $J = 7.3$  Hz, 3H).

### **(5-Cyano-7-methoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid**

To a solution of crude [5-cyano-8-methyl-1-propyl-7-(2,2,2-trichloro-acetimidoyloxymethyl)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid ethyl ester (45.7 mg, 0.089 mmol) in DCM/cyclohexanol (0.5 mL/0.5 mL) at 0°C was added methanol (14  $\mu$ L,



0.35 mmol) followed by trifluoromethanesulfonic acid (2 mL, 0.022 mmol). The cold bath was removed, and the solution stirred at ambient temperature for 20 minutes. The reaction was quenched by the addition of saturated  $\text{NaHCO}_3$  and extracted with EtOAc. The organic layer was washed with  $\text{H}_2\text{O}$  and brine, dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated. The crude material was chromatographed on  $\text{SiO}_2$  with 0.5% diethyl ether in DCM to 4% diethyl ether/DCM gradient to afford 13.3 mg (53%) of a colorless oil.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  9.67 (s, 1H), 7.40 (s, 1H), 4.55 (s, 2H), 4.21 (m, 2H), 4.05 (m, 1H), 3.91 (m, 1H), 3.34 (s, 3H), 3.06 (m, 3H), 2.91 (d,  $J = 16.4$  Hz, 1H), 2.53 (s, 3H), 2.11 (m, 1H), 1.95 (m, 1H), 1.29 (m, 5H), 0.88 (t,  $J = 7.3$  Hz, 3H).

The ethyl ester was saponified using 10% NaOH (aq., 150  $\mu\text{L}$ ) in methanol (2 mL) overnight to afford 8.8 mg (71%) of the carboxylic acid. ESI-MS  $m/z$  355 (M-H) $^-$ ;  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ )  $\delta$  10.71 (s, 1H), 7.38 (s, 1H), 4.58 (s, 2H), 4.03 (m, 2H), 3.39 (s, 3H), 3.03 (d,  $J = 14.1$  Hz, 1H), 2.98 (m, 2H), 2.85 (d,  $J = 14.1$  Hz, 1H), 2.56 (s, 3H), 2.04 (m, 2H), 1.42 (m, 1H), 1.01 (m, 1H), 0.87 (t,  $J = 7.3$  Hz, 3H).

Examples 3-19 were synthesized following the above mentioned procedure for example 1 using the intermediate [5-cyano-8-methyl-1-propyl-7-(2,2,2-trichloro-acetimidoyloxymethyl)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid ethyl ester and coupling it with the following alcohols: ethanol, 1-propanol, 2-propanol, cyclobutanol, cyclohexanol, cyclopropanemethanol, cyclobutanemethanol, cyclopentanemethanol, 2-butyne-1-ol, tetrahydro-4H-pyran-4-ol, (S)-3-hydroxytetrahydrofuran, (R)-3-hydroxytetrahydrofuran, benzyl alcohol, piperonyl alcohol, 2,4-dimethylbenzyl alcohol, 3-thiophenemethanol, and 2,4-dimethylthiazole-5-methanol. The resulting esters were hydrolyzed using 10% NaOH (aq) in ethanol (EtOH).

### Example 20

#### **(5-Cyano-8-methyl-7-phenoxyethyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid**

To a solution of (5-cyano-7-hydroxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid ethyl ester (42mg, 0.11 mmol) in THF (1.1 mL) was added triphenylphosphine (42 mg, 0.16 mmol) and phenol (15.3 mg, 0.16 mmol).

Diethylazodicarboxylate (DEAD, 26  $\mu\text{L}$ , 0.16 mmol) was added via syringe, and the reaction

was stirred for 1.5 hours, at which point additional portions of DEAD and triphenylphosphine were added. The reaction was stirred for an additional 1.5 hours. EtOAc and 10% NaOH were added, and the layers were separated. The organic layer was washed 10% NaOH (2 x), 3% HCl (1 x), saturated NaHCO<sub>3</sub> (1 x), and brine; dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated. Flash chromatography of the crude product on SiO<sub>2</sub> with 5% EtOAc/hexane to 15% EtOAc/hexane gradient afforded 16.3 mg (32%) of a colorless solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 9.76 (s, 1H), 7.50 (s, 1H), 7.3 (m, 3H), 6.98 (m, 2H), 5.13 (s, 2H), 4.26 (m, 2H), 4.12 (m, 1H), 3.95 (m, 1H), 3.08 (m, 2H), 3.01 (d, J = 17.0 Hz, 1H), 2.92 (d, J = 17.0 Hz, 1H), 2.56 (s, 3H), 2.04 (m, 1H), 1.97 (m, 1H), 1.3 (m, 5H), 0.89 (t, J = 7.3 Hz, 3H).

The ethyl ester was saponified using 10% NaOH (aq., 150 μL) in ethanol (1.5 mL) overnight to afford 15.7 mg (99%) of the carboxylic acid. ESI-MS m/z 417 (M-H)<sup>-</sup>; <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) δ 11.16 (br s, 1H), 7.58 (s, 1H), 7.32 (m, 2H), 7.05 (m, 2H), 6.96 (t, J = 7.3 Hz, 1H), 5.20 (s, 2H), 3.98 (m, 2H), 2.98 (d, J = 13.5 Hz, 1H), 2.88 (m, 2H), 2.75 (d, J = 13.5 Hz, 1H), 2.57 (s, 3H), 2.00 (m, 2H), 1.25 (m, 1H), 0.80 (m, 4H).

Example 21 was prepared using the above mentioned procedure for Example 20 using the intermediate (5-cyano-7-hydroxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid ethyl ester and coupling it with 3-fluorophenol. The resulting ester was hydrolyzed using 10% NaOH (aq) in EtOH.

## Example 22

**(5-Cyano-7-cyclopropylmethoxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid**

## 5-Bromo-2-methyl-1,3-dinitro-benzene

2,6-Dinitrotoluene (1.0 g, 5.5 mmol) was suspended in 4 mL con. H<sub>2</sub>SO<sub>4</sub>. 1,3-Dibromo-5,5-dimethylhydantoin (0.86 g, 3 mmol) was added portionwise to the mixture over 10 minutes. After a slight exotherm, the mixture went into solution briefly, and then a precipitate formed. The mixture was stirred at room temperature for 1 hour, and then the solid was filtered off. The

pale yellow solid was dried on a vacuum pump to give 1.33 g (92%) of the product. Purity 94% (GC); MS 260, 262 m/z (Br pattern);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  8.13 (s, 2H), 2.52 (s, 3H).

### 5-Bromo-2-methyl-3-nitro-phenylamine

5-Bromo-2-methyl-1,3-dinitro-benzene (1.0 g, 3.8 mmol) was mixed with EtOH (23 mL) and pyridine (1.56 mL, 19.0 mmol) and heated to reflux. A 20% solution of ammonium sulfide (3.89 g) in water was further diluted with 4 mL water then added via an addition funnel to the refluxing mixture over 1 hour. After the addition was complete, the reflux was continued for 2 hours. The reaction was cooled to room temperature, then poured onto a 1:1 mixture of water and ice (200 mL). The bright yellow precipitated solid was filtered off and dried on a vacuum pump to give 0.73 g (83%) of the product. Purity 100% (GC); MS 230, 232 m/z (Br pattern);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.29 (d,  $J = 1.7$  Hz, 1H), 6.99 (d,  $J = 1.7$  Hz, 1H), 3.97 (bs, 2H), 2.18 (s, 3H).

### 5-Bromo-2-methyl-3-nitro-phenol

5-Bromo-2-methyl-3-nitro-phenylamine (5.0 g, 21.5 mmol) was suspended in a solution of conc.  $\text{H}_2\text{SO}_4$  (6.4 mL) and water (21.2 mL). The mixture was cooled with an ice bath to  $\sim 0^\circ\text{C}$ , then a solution of sodium nitrite (1.66 g, 24.0 mmol) in water (6.4 mL) was added via an addition funnel at a rate to maintain the temperature below  $10^\circ\text{C}$ . This mixture was stirred at this temperature for 1 hour. The mixture was transferred via a plastic canula to a solution of conc.  $\text{H}_2\text{SO}_4$  (21.2 mL) and water (14.9 mL) heated at  $130\text{--}150^\circ\text{C}$ . The internal temperature of the mixture dropped to  $\sim 80^\circ\text{C}$  during the addition. The mixture was heated until the internal temperature returned to at least  $110^\circ\text{C}$ . The mixture was cooled slightly then poured on ice (200 mL), followed by an extraction with *t*-butylmethyl ether (3x100 mL). The organic layer was dried over  $\text{MgSO}_4$ , filtered, and solvent removed under reduced pressure to give 3.61 g (72%) of the product as a dark orange-red solid. Purity 98.5% (GC); MS 231, 233 m/z (Br pattern);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  10.82 (s, 1H), 7.53 (d,  $J = 1.7$  Hz, 1H), 7.25 (d,  $J = 1.7$  Hz, 1H), 2.17 (s, 3H).

### 1-Benzyloxy-5-bromo-2-methyl-3-nitro-benzene

To a slurry of NaH (767 mg, 32 mmol) in DMF (10 mL) was added a solution of 5-bromo-2-methyl-3-nitro-phenol (6.4 g, 27 mmol) in DMF (10 mL) at 0°C. After 15 minutes, the solution was treated with benzyl bromide (3.4 mL, 28 mmol) dropwise. The reaction mixture was warmed to room temperature. After 16 hours, the solution was carefully diluted with brine and extracted with diethyl ether (3x). The combined organics were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The crude material was purified by flash chromatography (SiO<sub>2</sub>, 6/1 hexanes/EtOAc) to give 5.4g (62%) of a tan-orange solid. GCMS M<sup>+</sup> 321 m/z. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.57 (s, 1H), 7.42-7.36 (m, 6H), 5.10 (s, 2H), 2.35 (s, 3H).

### **3-Benzyloxy-4-methyl-5-nitro-benzonitrile**

A solution of 1-benzyloxy-5-bromo-2-methyl-3-nitro-benzene (11 g, 34 mmol) in NMP (40 mL) was treated with CuCN (18g, 202 mmol) at room temperature. This solution was heated at 180°C for 1 hour. The solution was then cooled to room temperature, diluted with H<sub>2</sub>O and EtOAc and filtered through a Celite/silica gel pad. The filter cake was thoroughly rinsed with EtOAc. The filtrate was extracted with EtOAc and brine (2x), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The residual oil was diluted with diethyl ether and washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. Purification of the residual oil by flash chromatography (SiO<sub>2</sub>, 5/1 hexanes/EtOAc) followed by trituration with Et<sub>2</sub>O/hexanes provided 3.9 g (43%) of the benzonitrile as an off-white solid in addition to 3 g (37%) of the cyano-aniline as a light tan solid. GCMS M<sup>+</sup> 268 m/z. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.69 (s, 1H), 7.43-7.39 (m, 6H), 5.17 (s, 2H), 2.47 (s, 3H).

### **3-Amino-5-benzyloxy-4-methyl-benzonitrile**

A solution of 3-benzyloxy-4-methyl-5-nitro-benzonitrile (3.9 g, 14.5 mmol) in EtOH (40 mL) was treated with Fe powder (2.5 g) followed by acetic acid (4.0 mL) at room temperature. The solution was then heated at reflux for 2 hours. The solution was cooled to room temperature, filtered through a pad of Celite/silica gel and rinsed with EtOAc. The filtrate was concentrated *in vacuo*, extracted with EtOAc and saturated NaHCO<sub>3</sub> (2x), washed with brine,

dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to give 3.3 g (96%) of a light tan solid. GCMS M<sup>+</sup> 238 m/z. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.43-7.32 (m, 6H), 6.62 (s, 1H), 5.04 (s, 2H), 3.80 (bs, 2H), 2.11 (s, 3H).

### **3-Amino-5-benzyloxy-2-iodo-4-methyl-benzonitrile**

A solution of 3-amino-5-benzyloxy-4-methyl-benzonitrile (4.2 g, 22 mmol) in dichloroethane (40 mL) and methanol (20 mL) was treated with CaCO<sub>3</sub> (9 g, 90 mmol) followed by benzyltrimethylammonium dichloroiodate (9.2 g, 26 mmol) at room temperature. This solution was heated at reflux for 7 hours. The reaction was then cooled to room temperature, filtered through a Celite/silica gel pad and thoroughly rinsed with EtOAc. The filtrate was extracted (2x) with 10% NaHSO<sub>3</sub>, brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The residue was purified by flash chromatography (SiO<sub>2</sub>, 4/1 hexanes/EtOAc) followed by trituration with diethyl ether /hexanes to give 5.9 g (86%) of a light tan solid. GCMS M<sup>+</sup> 364 m/z. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.41-7.35 (m, 5H), 6.71 (s, 1H), 5.04 (s, 2H), 4.36 (bs, 2H), 2.20 (s, 3H).

### **6-Benzyloxy-3-(2-hydroxy-ethyl)-7-methyl-2-triethylsilanyl-1H-indole-4-carbonitrile**

A solution of 4-triethylsilanyl-but-3-yn-1-ol (913 mg, 4.9 mmol) in DMF (2 mL) was treated with 3-amino-5-benzyloxy-2-iodo-4-methyl-benzonitrile (497 mg, 1.4 mmol) in DMF (2 mL) followed by tetrabutylammonium chloride (401 mg, 1.4 mmol), triphenylphosphine (115 mg, 0.43 mmol), palladium acetate (90 mg, 0.40 mmol), and diisopropylethylamine (1.0 mL, 5.7 mmol) at room temperature. This solution was heated at 85°C for 3.5 hours. After cooling the reaction to room temperature, the reaction mixture was filtered through a Celite/silica gel pad. The filter pad was thoroughly washed with EtOAc. The filtrate was extracted with brine (2x), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The residue was diluted with diethyl ether and washed with brine (2x), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The crude oil was purified by flash chromatography (SiO<sub>2</sub>, 4/1-1/1 hexanes/EtOAc) to give 422 mg (72%) of an orange oil that solidified upon standing. LCMS (M-H) 419 m/z. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.99 (s, 1H), 7.47-7.34 (m, 5H), 7.17 (s, 1H), 5.12 (s, 2H), 3.95-3.91 (m, 2H), 3.33-3.28 (m, 2H), 2.47(s, 3H), 2.12-2.09 (m, 1H), 1.05-0.92 (m, 15H).

**(±)-(7-Benzyloxy-5-cyano-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl)-acetic acid ethyl ester**

A solution of 6-benzyloxy-3-(2-hydroxy-ethyl)-7-methyl-2-triethylsilyl-1*H*-indole-4-carbonitrile (242 mg, 0.58 mmol) in DCM (4 mL) was treated with BF<sub>3</sub>-OEt<sub>2</sub> (0.08 mL, 0.64 mmol) at room temperature. After 0.5 hours, additional BF<sub>3</sub>-OEt<sub>2</sub> (0.10 mL) was added. After 0.5 hours, ethyl butyrlacetate (0.12 mL, 0.75 mmol) was added. After 3 hours, the reaction was diluted with brine, extracted with EtOAc (3x), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The residue was triturated with diethyl ether /hexanes to give 130 mg (50%) of the pyran as an off-white solid. LCMS (M-H) 445 m/z. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 9.44 (s, 1H), 7.45-7.33 (m, 5H), 7.10 (s, 1H), 5.11 (s, 2H), 4.29-4.11 (m, 2H), 4.08-4.02 (m, 1H), 3.95-3.79 (m, 1H), 3.06-3.02 (m, 2H), 2.96 (q, *J*<sub>AB</sub> = 18 Hz, 2H), 2.43 (s, 3H), 2.12-1.83 (m, 2H), 1.42-1.33 (m, 1H), 1.29 (t, *J* = 6.9 Hz, 3H), 1.23-1.14 (m, 1H), 0.88 (t, *J* = 7.2 Hz, 3H).

**(±)-(5-Cyano-7-hydroxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl)-acetic acid ethyl ester**

A solution of (±)-(7-benzyloxy-5-cyano-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl)-acetic acid ethyl ester (143 mg, 0.32 mmol) in EtOAc (10 mL) and MeOH (10 mL) was treated with Pd(OH)<sub>2</sub> (20 mg) and subjected to H<sub>2</sub> using Parr shaker. After 1 hour, the solution was filtered through a Celite plug. The filter pad was thoroughly rinsed with EtOAc. The filtrate was concentrated, and the residue was then triturated with diethyl ether /hexanes to give the phenol as an off-white solid (69 mg, 61%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 9.45 (s, 1H), 6.94 (s, 1H), 4.65 (s, 1H), 4.26-4.11 (m, 2H), 4.07-4.00 (m, 1H), 3.94-3.86 (m, 1H), 3.04-3.01 (m, 2H), 2.96 (q, *J*<sub>AB</sub> = 17 Hz, 2H), 2.41 (s, 3H), 2.12-2.01 (m, 1H), 1.96-1.86 (m, 1H), 1.43-1.33 (m, 1H), 1.29 (t, *J* = 7.5 Hz, 3H), 1.23-1.17 (m, 1H), 0.88 (t, *J* = 7.5 Hz, 3H).

**(5-Cyano-7-cyclopropylmethoxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl)-acetic acid**

To a solution of (5-cyano-7-hydroxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid ethyl ester (36.8 mg, 0.10 mmol) in THF (2 mL) was added triphenyl phosphine (52 mg, 0.20 mmol) and cyclopropanemethanol (16  $\mu$ L, 0.2 mmol).

Diisopropylazodicarboxylate (DIAD, 39  $\mu$ L, 0.2 mmol) was added. The reaction was stirred for 20 minutes, then quenched with H<sub>2</sub>O and diluted with EtOAc. The organic layer was washed with H<sub>2</sub>O (1x) and brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated. Flash chromatography on SiO<sub>2</sub> with 8% EtOAc/hexane afforded 29.5 mg (70%) of the product as a white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  9.43 (br s, 1H), 7.03 (s, 1H), 4.21 (m, 2H), 4.07 (m, 1H), 3.9 (m, 3H), 3.02 (m, 3H), 2.90 (d, J = 16.4 Hz), 1H), 2.43 (s, 3H), 2.03 (m, 1H), 1.96 (m, 1H), 1.3 (m, 5H), 0.88 (t, J = 7.3 Hz), 0.64 (m, 2H), 0.37 (m, 2H).

To a solution of the ethyl ester (29.4 mg, 0.072 mmol) in EtOH (2 mL) and THF (0.15 mL) was added 10% NaOH (aq., 0.2 mL). After stirring for 17.25 hours, the solution was acidified with 3% HCl and extracted with EtOAc. The organic layer was washed with H<sub>2</sub>O and brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to afford 26.9 mg (98%) of the title compound as a white solid. ESI-MS m/z 383 (MH)<sup>+</sup>. <sup>1</sup>H NMR (CD<sub>3</sub>OD)  $\delta$  7.07 (s, 1H), 4.05 (m, 2H), 3.87 (d, J = 6.4 Hz), 2.99 (m, 3H), 2.83 (d, J = 14.1 Hz, 1H), 2.43 (s, 3H), 2.02 (m, 2H), 1.40 (m, 2H), 1.02 (m, 1H), 0.86 (t, J = 7.3 Hz, 3H), 0.61 (m, 2H), 0.36 (m, 2H).

### Example 23

**(R)-[5-Cyano-8-methyl-7-(5-methyl-isoxazol-3-ylmethoxy)-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid**

**(R)-(5-Cyano-7-hydroxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid ethyl ester**

Preparative chiral HPLC using CHIRALPAK AS (250 x 20mm) and 20% 2-propanol in heptane as eluant gave the (R) and (S) enantiomers of (5-cyano-7-hydroxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid ethyl ester as white solids. Analytical chiral HPLC HP 100 with CHIRALPAK AS, 250 x 4.6 mm, 2-propanol/heptane (20/80), 1.0 mL/min, 235 nm UV detection; t<sub>R</sub> = 5.70 min (R enantiomer), 12.74 min (S enantiomer).

**(R)-[5-Cyano-8-methyl-7-(5-methyl-isoxazol-3-ylmethoxy)-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid**

To a solution of (R)-(5-cyano-7-hydroxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid ethyl ester (0.17 g, 0.47 mmol) in dry DMF (3mL) was added milled potassium carbonate (71 mg, 0.52 mmol). This was stirred for 15 minutes, at which point 3-bromomethyl-5-methyl-isoxazole (91 mg, 0.52 mmol) was added. The reaction was stirred for 16 hours at ambient temperature, diluted with H<sub>2</sub>O and extracted with EtOAc. The organic layer was washed several times with H<sub>2</sub>O and once with brine, dried (MgSO<sub>4</sub>), and concentrated. The residue was flash chromatographed on SiO<sub>2</sub> using 20% EtOAc in hexanes resulting in 185 mg (88% yield) of the product as an oil which crystallized upon standing.

To a solution of (R)-[5-cyano-8-methyl-7-(5-methyl-isoxazol-3-ylmethoxy)-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid ethyl ester (185 mg, 0.40 mmol) in ethanol (10mL) was added 2M sodium hydroxide (0.4mL). After stirring for 36 hours, the reaction was concentrated in vacuo. The residue was taken up in water and extracted with EtOAc. The aqueous layer was acidified and re-extracted with EtOAc. The combined organic layers were dried (MgSO<sub>4</sub>) and concentrated to afford 150 mg (92%) of the title compound. ESI-MS m/z 424 (MH)<sup>+</sup>. <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 7.20 (s, 1H), 6.25 (s, 1H), 5.15 (s, 2H), 3.0 (m, 3H), 2.83 (d, J = 14.1 Hz, 1H), 2.43 (s, 3H), 2.42 (s, 3H), 2.04 (m, 2H), 1.40 (m, 1H), 1.01 (m, 1H), 0.86 (t, J = 7.3 Hz, 3H).

**Example 24**

**[5-Cyano-8-methyl-1-propyl-7-(pyridin-4-ylmethoxy)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid**

To a solution of (R)-(5-cyano-7-hydroxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid ethyl ester (140 mg, 0.39 mmol) in THF (6 mL) was added 4-pyridylcarbinol (87 mg, 0.80 mmol) and triphenylphosphine (207 mg, 0.80 mmol). After cooling to 0°C, DIAD (157 μL, 0.80 mmol) was added. The cold bath was removed, and the solution was stirred at ambient temperature for 2.5 hours. The reaction was quenched with water and extracted with EtOAc. The organic layer was washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated. The residue was dissolved in diethyl ether, and 4N HCl in dioxane (ca. 1mL) was



added. The precipitated solids were collected by filtration, washed with ether, and partitioned between EtOAc and 5% NaHCO<sub>3</sub>. The aqueous layer was extracted with EtOAc. The combined organic layers were washed with H<sub>2</sub>O and brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to afford 95.9 mg (54%) of a pale yellow solid.

To a solution of the ethyl ester (95 mg, 0.21 mmol) in EtOH (5 mL) was added 10% NaOH (aq., 0.5 mL). After stirring for 18 hours, H<sub>2</sub>O was added and the pH brought to pH 6-7 with 3% HCl. The mixture was extracted with EtOAc (3 x). The combined organic layers were washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to afford 67.3 mg (76%) of the title compound as a pale yellow solid. ESI-MS *m/z* 420 (MH)<sup>+</sup>. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) δ 10.97 (s, 1H), 8.60 (m, 2H), 7.47 (m, 2H), 7.28 (s, 1H), 5.25 (s, 2H), 3.93 (m, 2H), 2.94 (d, *J* = 14.1 Hz, 1H), 2.83 (m, 2H), 2.72 (d, *J* = 14.1 Hz, 1H), 2.47 (s, 3H), 1.99 (m, 2H), 1.20 (m, 2H), 0.80 (m, 3H).

#### Example 25

**(R)-[5-Cyano-7-(1,5-dimethyl-1H-pyrazol-3-ylmethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid**

**(±)-(5-Cyano-7-hydroxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid**

To a solution of (±)-(5-cyano-7-hydroxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid ethyl ester (3.0 g, 8.43 mmol) in ethanol (120mL) was added 2M sodium hydroxide (10mL). After stirring for 20 hours, the reaction mixture was concentrated, and the resulting residue was taken up in water. The aqueous solution was acidified, and the solids collected by filtration and washed with water. The wet solids were dried overnight in a vacuum oven at 40°C to afford 2.7g (96% yield) of the product. ESI-MS *m/z* 329 (MH)<sup>+</sup>. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) δ 11.96 (s, 1H), 10.71 (s, 1H), 9.35 (s, 1H), 6.93 (s, 1H), 3.95 (m, 2H), 2.93 (d, *J* = 13.5 Hz, 1H), 2.80 (m, 2H), 2.71 (d, *J* = 13.5 Hz, 1H), 2.33 (s, 3H), 1.96 (m, 2H), 1.28 (m, 1H), 0.80 (m, 4H).

**(R)-(5-Cyano-7-hydroxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid methyl ester**

To a solution of (+/-)(5-cyano-7-hydroxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid ( 2.7 g, 8.22 mmol) in methanol (50mL) at 60°C was added a solution of (-) quinine (2.9 g, 8.94 mmol) in methanol (40mL). After 30 minutes at 60°C, a white solid precipitated. The solution was allowed to stir for 20 hours at 60°C. After cooling to room temperature, the solids were recovered by filtration and washed with methanol to afford 2.3 g of the quinine salt of (R)-(5-cyano-7-hydroxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid. The solids were partitioned between 1M HCl (50mL) and EtOAc (50mL). The organic layer was washed with brine, dried (MgSO<sub>4</sub>), and concentrated to afford (R)-(5-cyano-7-hydroxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid as a white solid. The solids were taken up in toluene (30mL) and trimethylorthoacetate (60mL), and the solution was heated at 100°C for 2 hours. The reaction was concentrated in vacuo, and the resulting residue was chromatographed on SiO<sub>2</sub> eluting with 30% EtOAc in hexanes to afford 1.25g (46%) of the methyl ester whose chiral purity was determined to be 98.5% (97% ee, Chiral HPLC HP 100 with CHIRALPAK AS, 250 x 4.6 mm, 2-propanol/heptane (20/80), 1.0 mL/min, 235 nm UV detection; t<sub>R</sub> = 6.35 min (R enantiomer), 8.53 min (S enantiomer)). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 9.40 (s, 1H), 6.94 (s, 1H), 4.69 (s, 1H), 4.02 (m, 1H), 3.92 (m, 1H), 3.75 (s, 3H), 2.94-3.06 (m, 2H), 2.41 (s, 3H), 1.91- 2.04 (m, 2H), 1.26-1.53 (m, 2H), 0.88 (t, J = 7.0 Hz, 3H).

**[5-Cyano-7-(1,5-dimethyl-1H-pyrazol-3-ylmethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid**

To a solution of 5-cyano-7-hydroxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid methyl ester (50 mg, 0.15mmoles) in DCM (1 mL) was added (1,5-dimethyl-1H-pyrazol-3-yl)-methanol (20 mg, 0.17mmoles) and triphenylphosphine (95mg, 0.37 mmol). To this was added 1,1'-(azodicarbonyl)dipiperidine (ADDP, 92 mg, 0.37mmoles). The reaction was stirred for 1.5 hours, diluted with EtOAc and washed twice with H<sub>2</sub>O and brine.

The EtOAc layer was dried (MgSO<sub>4</sub>) and concentrated. Flash chromatography on SiO<sub>2</sub> eluting with 1:1 hexane/EtOAc yielded 53 mg (80%) of a yellow oil that crystallized upon standing.

To a solution of the methyl ester (53 mg, 0.12 mmol) in EtOH (5 mL) was added 2N NaOH (1 mL). The reaction was stirred at 40°C for 1 hour, cooled, and concentrated *in vacuo*. The residue was taken up in H<sub>2</sub>O, acidified, and extracted with EtOAc. The combined organic layers were washed with brine, dried (MgSO<sub>4</sub>), and concentrated. Trituration of the solids with DCM/hexane afforded 36 mg (71%) of the acid as an off-white solid. ESI-ms *m/z* 437 (MH)<sup>+</sup>. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) δ 11.98 (s, 1H), 10.87 (s, 1H), 7.32 (s, 1H), 6.10 (s, 1H), 4.98 (s, 2H), 3.95 (m, 2H), 3.69 (s, 3H), 2.93 (d, *J* = 13.5 Hz, 1H), 2.82 (m, 2H), 2.70 (d, *J* = 13.5 Hz, 1H), 2.35 (s, 3H), 2.23 (s, 3H), 1.96 (m, 2H), 1.16 (m, 1H), 0.82 (m, 4H).

### Example 26

**(R)-[5-Cyano-7-(2-isopropoxy-ethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid**

#### **2-(2-Bromoethoxy)propane**

To a 2 neck flask equipped with a reflux condenser, magnetic stir bar and a septum was added 2-isopropoxyethanol (2 mL, 17.34 mmol). Phosphorous tribromide (0.52 mL, 5.45 mmol) was added dropwise via syringe. The reaction temperature increased and a slight reflux occurred. The reaction was stirred at room temperature under argon for 24 hours, quenched with water and extracted with hexane. The organic layer was dried over magnesium sulfate and concentrated to provide 1 g (34%) of the product as a colorless oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 3.78 (t, 2H); 3.61 (m, 1 H); 3.42 (t, 2H); 1.18 (d, 6H).

**(R)-[5-Cyano-7-(2-isopropoxy-ethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid**

To a solution of (R)-(5-cyano-7-hydroxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl)-acetic acid methyl ester (125 mg, 0.365 mmol) and 2-(2-bromoethoxy)propane (0.07 mL, 0.539 mmol) in anhydrous N,N-dimethylformamide (2.5 mL)

was added cesium carbonate (250 mg, 0.767 mmol) and a catalytic amount of sodium iodide. The reaction was heated at 60°C for 1 hour, cooled to room temperature, quenched with water and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and concentrated. Purification by column chromatography on silica gel using 3/1 hexane/ethyl acetate as eluent provided 52 mg (33%) of the product as a white solid. ESI-MS  $m/z$  429 (M+H)<sup>+</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  9.36 (s, 1H); 7.07 (s, 1H); 4.14 (t, 2H); 4.05 (m, 1H); 3.90 (m, 1H); 3.78 (t, 2H); 3.74 (s, 3H); 3.68 (m, 1H); 3.03 (m, 3H); 2.91 (d, 1H), 2.42 (s, 3H); 2.03 (m, 1H); 1.96 (m, 1H); 1.39 (m, 1H), 1.22 (d, 6H); 1.17 (m, 1H); 0.94 (t, 3H).

To a solution of the methyl ester (48 mg, 0.112 mmol) in ethanol (2 mL) was added 10% aqueous sodium hydroxide solution (0.2 mL, excess). After stirring at room temperature for 24 hours, the solvent was removed by rotary evaporation. The residue was dissolved in water and acidified with 1N HCl. The solid precipitate was collected by suction filtration, washed with water and dried to provide 43 mg (93%) of the product as a white solid. ESI-MS  $m/z$  413.2 (M-H)<sup>-</sup>. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO)  $\delta$  11.98 (s, 1H); 10.90 (s, 1H); 7.20 (s, 1H); 4.11 (t, 2H); 3.89 (m, 2H); 3.69-3.58 (m, 3H); 2.90 (d, 1H), 2.81 (m, 2H); 2.68 (d, 1H); 2.37 (s, 3H); 1.96 (m, 2H); 1.28 (m, 1H); 1.10 (d, 6H); 0.76 (m, 4H).

### Example 27

#### **(R)-[5-Cyano-7-(3-methoxy-propoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid**

To a solution of (R)-(5-cyano-7-hydroxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl)-acetic acid methyl ester (100 mg, 0.292 mmol) and 3-methoxypropyl bromide (67 mg, 0.438 mmol) in anhydrous N,N-dimethylformamide (2 mL) was added cesium carbonate (143 mg, 0.438 mmol) and a catalytic amount of sodium iodide. The reaction was heated at 60°C for 24 hours, cooled to room temperature, quenched with water and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and concentrated. Purification by column chromatography on silica gel using 2/1 hexane/ethyl acetate as eluent provided the product as a white solid. ESI-MS  $m/z$  415 (M+H)<sup>+</sup>.

To a solution of the methyl ester in ethanol (2 mL) was added 10% aqueous sodium hydroxide solution (0.2 mL, excess). After stirring at room temperature for 24 hours, the solvent

was removed by rotary evaporation. The residue was dissolved in water and acidified with 1N HCl. The solid precipitate was collected by suction filtration, washed with water and dried to provide 33 mg (28% over two steps) of the product as a white solid. ESI-MS  $m/z$  399 (M-H)<sup>-</sup>. <sup>1</sup>H NMR (DMSO)  $\delta$  11.99 (s, 1H); 10.87 (s, 1H); 7.17 (s, 1H); 4.05 (t, 2H); 3.91 (m, 2H); 3.49 (t, 2H); 3.24 (s, 3H); 2.89 (d, 1H); 2.81 (m, 2H); 2.72 (d, 1H); 2.36 (s, 3H); 1.94 (m, 4H); 0.77 (m, 4H).

### Example 28

**(1R, 2'R)-[5-Cyano-7-(2-methoxy-propoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid**

#### **(R)-2-methoxypropan-1-ol**

To a 2 neck flask fitted with a reflux condenser, an argon inlet and a septum was added lithium aluminum hydride (442 mg, 11.64 mmol) and diethyl ether (5 mL). The suspension was placed under argon, and a solution of ethyl (R)-(-)-2-methoxypropionate (2 g, 15.13 mmol) in diethyl ether (5 mL) was added dropwise via syringe. The mixture was heated at reflux for 1 hour, cooled to room temperature, and a solution of potassium hydroxide (1.7 g, 30.30 mmol) in water (4 mL) was added. The ether layer was decanted, and the aqueous mixture was washed twice more with ether followed by decanting. The ether washes were combined, dried over magnesium sulfate and evaporated in a cold water bath under reduced pressure. Distillation of the mixture gave 818 mg (60 %) of the product as a colorless oil, b.p = 130°C. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  3.59 (m, 1H), 3.40 (s, 3H); 2.19 (br s, 1H); 1.39 (d, 2H); 1.11 (d, 3H).

**(1R, 2'R)-[5-Cyano-7-(2-methoxy-propoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid**

To a solution of (R)-(5-cyano-7-hydroxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid methyl ester (100 mg, 0.292 mmol), triphenylphosphine (115 mg, 0.438 mmol) and 1,1'-(azocarbonyl)dipiperidine (110 mg, 0.438 mmol) in anhydrous dichloromethane (2 mL) was added a solution of 2-methoxypropan-1-ol (39 mg, 0.438 mmol)

The reaction was stirred at room temperature under argon for 48 hours, and the solvent removed by rotary evaporation. Purification by column chromatography on silica gel using 2/1 hexane/ethyl acetate as eluent provided the product as a white solid. ESI-MS  $m/z$  415 (M+H)<sup>+</sup>

To a solution of the methyl ester in ethanol (2 mL) was added 10% aqueous sodium hydroxide solution (0.2 mL, excess). After stirring at room temperature for 24 hours, the solvent was removed by rotary evaporation. The residue was dissolved in water and acidified with 1N HCl. The solid precipitate was collected by suction filtration, washed with water and dried to provide 27 mg (23% over two steps) of the product as a white solid. ESI-MS  $m/z$  399 (M-H)<sup>-</sup>. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO)  $\delta$  11.99 (s, 1H), 10.88 (s, 1H); 7.21 (s, 1H); 3.99-3.88 (m, 4H); 3.67(m,1H); 3.34 (s, 3H); 2.94 (d, 1H); 2.83 (m, 2H); 2.71 (d, 1H); 2.39 (s, 3H), 1.97 (m, 2H); 1.29 (m, 1H); 1.20 (d, 3H); 0.76 (m, 4H).

#### Example 29

**[5-Cyano-8-methyl-7-(5-methyl-[1,3,4]thiadiazol-2-ylmethoxy)-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid**

**Chloro-acetic acid N<sup>1</sup>-acetyl-hydrazide.**

In a 50-mL round bottom flask, 1.07 g (13.0 mmol) of acetic hydrazide was dissolved in water (5 mL) and combined with Na<sub>2</sub>CO<sub>3</sub> (0.814 g, 7.69 mmol). The resulting mixture was first cooled in an ice-bath and then chloroacetyl chloride (1.1 mL, 13.8 mmol) was added dropwise over 10 minutes with stirring. After all of the chloroacetyl chloride had been added, the reaction was allowed to warm to room temperature and to continue to stir for 1.5 hours. The product was filtered out and dried *in vacuo* to afford 0.202 g (10%) of clean chloro-acetic acid N<sup>1</sup>-acetyl-hydrazide: <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  10.17 (s, 1H), 9.93 (s, 1H), 4.11 (s, 2H), 1.86 (s, 3H).

#### 2-Chloromethyl-5-methyl-[1,2,4]thiadiazole

In a 250-mL round bottom flask, chloro-acetic acid N<sup>1</sup>-acetyl-hydrazide (0.400 g, 2.66 mmol) was combined with THF (40 mL) and phosphorus pentasulfide (P<sub>2</sub>S<sub>5</sub>) (1.83 g, 4.12 mmol). The resulting mixture was heated at reflux for 2 hours and then cooled to room

temperature. Once at room temperature, 5% Na<sub>2</sub>CO<sub>3</sub> (16 mL) and excess diethyl ether were added; the resulting suspension was filtered. The organic layer was separated, dried over MgSO<sub>4</sub>, filtered, and concentrated *in vacuo* to afford 0.160 g (41%) of 2-chloromethyl-5-methyl-[1,2,4]thiadiazole: <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 4.91 (s, 2H), 2.80 (s, 3H).

**(R)-[5-Cyano-8-methyl-7-(5-methyl-[1,3,4]thiadiazol-2-ylmethoxy)-1-propyl-1,3,4,9-tetrahydro-pyrano{3,4-*b*}indol-1-yl]-acetic acid methyl ester**

In a 50-mL round bottom flask and under argon, (R)- (5-cyano-7-hydroxy-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indole-1-yl)-acetic acid methyl ester (0.100 g, 0.292 mmol) was combined with 2-chloromethyl-5-methyl-[1,2,4]thiadiazole (0.108 g, 0.730 mmol) and anhydrous NMP (4 mL). With stirring, K<sub>2</sub>CO<sub>3</sub> (0.101 g, 0.730 mmol) and KI (0.012 g, 0.073 mmol) were added. The resulting mixture was allowed to continue stirring at room temperature for 24 hours. The reaction mixture was then poured onto water, and the product was extracted with EtOAc. The organic layer was washed twice with ice-water, once with brine, dried over MgSO<sub>4</sub>, filtered, and concentrated *in vacuo* to yield an oil that was then purified by flash chromatography to afford 0.084 g (63%) of (R)-[5-cyano-8-methyl-7-(5-methyl-[1,3,4]thiadiazol-2-ylmethoxy)-1-propyl-1,3,4,9-tetrahydro-pyrano{3,4-*b*}indol-1-yl]-acetic acid methyl ester: <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 9.52 (s, 1H), 7.11 (s, 1H), 5.46 (s, 2H), 4.07 (m, 1H), 3.94 (m, 1H), 3.75 (s, 3H), 3.06 (m, 4H), 2.81 (s, 3H), 2.34 (s, 3H), 2.11 (m, 2H), 1.43 (m, 1H), 1.28 (m, 1H), 0.90 (t, J = 7.3 Hz, 3H).

**(R)-[5-Cyano-8-methyl-7-(5-methyl-[1,3,4]thiadiazol-2-ylmethoxy)-1-propyl-1,3,4,9-tetrahydro-pyrano{3,4-*b*}indol-1-yl]-acetic acid**

In a 25-mL round bottom flask, (R)-[5-cyano-8-methyl-7-(5-methyl-[1,3,4]thiadiazol-2-ylmethoxy)-1-propyl-1,3,4,9-tetrahydro-pyrano{3,4-*b*}indol-1-yl]-acetic acid methyl ester (0.084 g, 0.185 mmol) was combined with EtOH (8 mL) and 10% NaOH (0.38 mL). The resulting solution was allowed to stir for 24 hours at room temperature. The reaction solution was concentrated *in vacuo* to yield a residue that was then taken up in water and washed once with diethyl ether. The aqueous layer was then made acidic with 4M HCl, and the precipitated

product was extracted with EtOAc, dried over MgSO<sub>4</sub>, and concentrated *in vacuo* to afford 0.069 g (85%) of (R)-[5-cyano-8-methyl-7-(5-methyl-[1,3,4]thiadiazol-2-ylmethoxy)-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid: Purity 98% (HPLC); MS (M-H)<sup>+</sup> 439.1 m/z; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 9.53 (s, 1H), 7.03 (s, 1H), 5.46 (d, J = 2.9Hz, 2H), 4.17 (m, 2H), 3.12 (m, 4H), 2.83 (s, 3H), 2.15 (s, 3H), 2.13 (m, 2H), 1.49 (m, 1H), 1.29 (m, 1H), 0.918 9 (t, J = 7.3Hz, 3H).

### Example 30

**(R)-[5-Cyano-7-(5-dimethylamino-[1,2,4]thiadiazol-3-ylmethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid**

#### **5-(N, N-Dimethylamino)-3-(chloromethyl)-1,2,4-thiadiazole**

5-Chloro-3-(chloromethyl)-1,2,4-thiadiazole (0.2 g, 1.2 mmol) was dissolved in dry THF (2 mL) followed by the addition of a 2M solution of dimethylamine in THF (1.18 mL, 2.4 mmol). The reaction was stirred at room temperature overnight. The precipitated solid was filtered off, and the solvent removed under reduced pressure. The residue was taken up in ether, and washed with water and sat. NaCl. The organic layer was dried over MgSO<sub>4</sub>, filtered, and the solvent was removed under reduced pressure to give an oil (0.210 g) in quantitative yield. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 4.51 (s, 2H), 3.16 (s, 6H).

**(R)-[5-Cyano-7-(5-dimethylamino-[1,2,4]thiadiazol-3-ylmethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid methyl ester**

(R)- (5-Cyano-7-hydroxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid methyl ester (0.1 g, 0.29 mmol) was dissolved in dry NMP (3 mL), followed by the addition of 5-(N, N-dimethylamino)-3-(chloromethyl)-1,2,4-thiadiazole (0.124 g, 0.70 mmol), milled potassium carbonate (0.096 g, 0.70 mmol) and potassium iodide (0.011 g, 0.070 mmol). The reaction was stirred at room temperature overnight. EtOAc (50 mL) was added to the reaction, and it was washed several times with ice water and finally with sat. NaCl. The organic layer was dried over MgSO<sub>4</sub>, filtered, and the solvent was removed under reduced pressure to give a residue that was purified by silica gel flash chromatography to give 0.104 g (74% yield)



product as a white solid. Purity 99.3% (HPLC); MS (M-H)<sup>-</sup> 482.2 m/z; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 9.38 (s, 1H), 7.18 (s, 1H), 5.11 (s, 2H), 4.05 (m, 1H), 3.90 (m, 1H), 3.75 (s, 3H), 3.18 (s, 6H), 3.03 (m, 3H), 2.92 (d, J = 16.4 Hz, 1H), 2.48 (s, 3H), 2.06 (m, 1H), 1.92 (m, 1H), 1.38 (m, 1H), 1.19 (m, 1H), 0.89 (t, J = 7.3 Hz, 3H).

**(R)-[5-Cyano-7-(5-dimethylamino-[1,2,4]thiadiazol-3-ylmethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid**

(R)-[5-Cyano-7-(5-dimethylamino-[1,2,4]thiadiazol-3-ylmethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid methyl ester (0.100 g, 0.207 mmol) was dissolved in EtOH (4 mL), followed by the addition of a 10% sodium hydroxide solution (0.41 mL). The reaction was stirred overnight at room temperature, then the solvent was removed under reduced pressure. The residue was dissolved in water (50 mL) and washed with ether (2X25 mL). The aqueous layer was acidified with 1M HCl to about pH 4, then extracted with EtOAc. The organic layer was dried over MgSO<sub>4</sub>, filtered, and solvent removed under reduced pressure to give the product (0.09 g, 93% yield) as an off white solid. Purity 97.5% (HPLC); MS (M+H)<sup>+</sup> 470.1 m/z; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 11.98 (s, 1H), 10.91 (s, 1H), 7.31 (s, 1H), 5.09 (s, 2H), 3.92 (m, 2H), 3.10 (s, 6H), 2.93 (d, J = 13.5 Hz, 1H), 2.82 (m, 2H), 2.71 (d, J = 13.5 Hz, 1H), 2.41 (s, 3H), 1.97 (m, 2H), 1.30 (m, 1H), 0.78 (m, 4H).

**Example 31**

**5-Cyano-7-(2-methoxy-ethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4,-b]indole-1-carboxylic acid**

**5-Bromo-2-methyl-3-nitrobenzoic acid**

A 5000 mL three neck RBF equipped with an overhead stirrer was charged with 2-methyl-3-nitro benzoic acid (150.0 g, 0.82 mol) and concentrated H<sub>2</sub>SO<sub>4</sub> (600 mL). To this solution was added 1,3-dibromo-5,5-dimethylhydantoin (130.7 g, 0.455 mol) over ten minutes with vigorous stirring. The reaction was vigorously stirred at ambient temperature for 5 hours. The reaction mixture was then added to H<sub>2</sub>O (4000 mL) chilled in an ice bath over 30 minutes.

This mixture was then filtered, the solids washed twice with H<sub>2</sub>O and further dried under vacuum to yield 217.7 g (99.8%) of an off-white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.18 (s, 1H), 7.95 (s, 1H), 2.59 (s, 3H).

### **5-Bromo-2-methyl-3-nitrobenzoic acid benzyl ester**

A 5000 mL three neck RBF equipped with an overhead stirrer and thermometer was charged with 5-bromo-2-methyl-3-nitro benzoic acid (116.2 g, 0.45 mol), THF (1L), and benzyl bromide (84.90 mL, 0.715 mol). To the stirring solution was added diisopropyl ethylamine (78 mL, 0.450 mol). The reaction was then brought to reflux. After 5.5 hours at reflux, the reaction was cooled to 40°C and pyrrolidine (83 mL, 1.00 mol) was added. The reaction was stirred 10 minutes at 40°C, and then allowed to cool to ambient temperature over approximately 20 minutes. The reaction was diluted with EtOAc and washed 2x 3% HCl, 1x H<sub>2</sub>O, 1x saturated NaCl, dried (MgSO<sub>4</sub>), and concentrated. The brown oil was taken up in DCM and slurried with approximately 300 g of silica gel. The slurry was filtered, solids washed with EtOAc, and the filtrate was concentrated to afford a yellow oil. Upon trituration in hexanes, 244.9 g (83.8%) of off white powder was obtained. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.10 (s, 1H), 7.96 (s, 1H), 7.41 (m, 5H), 5.37 (s, 2H), 2.55 (s, 3H).

### **3-Amino-5-bromo-2-methylbenzoic acid benzyl ester**

A 1000 mL RBF equipped with an overhead stirrer was charged with of 5-bromo-2-methyl-3-nitro benzoic acid benzyl ester (75.0 g, 0.214 mol) and H<sub>2</sub>O (200mL). While stirring vigorously, Fe powder (~325 mesh, 47.81g, 0.857 mol) and NH<sub>4</sub>Cl (13.88g, 0.257 mol) were added. The mixture was heated to reflux for 4.5 hours. The reaction was then allowed to cool to room temperature, diluted with EtOAc, and filtered through a pad of Celite. The filtrate was washed with H<sub>2</sub>O (2x) and saturated NaCl (1x), dried (MgSO<sub>4</sub>), and concentrated. Upon trituration in hexanes, 63.2 g (92.1%) of off-white powder was obtained. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.38 (m, 6H), 6.90 (s, 1H), 5.31 (s, 2H), 3.76 (br s, 2H), 2.25 (s, 3H).

### **5-Bromo-3-hydrazino-2-methyl benzoic acid benzyl ester hydrochloride**

A 1000 mL three neck RBF equipped with an overhead stirrer was charged with 3-amino-5-bromo-2-methyl benzoic acid benzyl ester (25.0 g, 0.078 mol), H<sub>2</sub>O (150mL), and concentrated HCl (150mL). This mixture was stirred vigorously while cooling to -10°C in a MeOH/ice bath. A solution of NaNO<sub>2</sub> (8.08 g, 0.120 mol) in H<sub>2</sub>O (150mL) was cooled to -10°C and then added to the reaction mixture dropwise over 15 minutes. The reaction was vigorously stirred at -10°C for 1.5 hours, then a -10°C solution of SnCl<sub>2</sub>·2H<sub>2</sub>O (73.99 g, 0.330 mol) in concentrated HCl (150mL) was added to the reaction mixture dropwise over 20 minutes. This was allowed to react for 1.5 hours at -10°C with very vigorous stirring. The reaction mixture was then added to 6N NaOH (600mL) and extracted with EtOAc. The organic layer was separated and washed with saturated NaCl, dried over MgSO<sub>4</sub>, and concentrated. The yellow solid was taken up in THF (100mL), diluted with 4N HCl dioxane (30 mL), and then the solvents were removed under vacuum. Trituration with DCM/hexanes yielded 27.1 g (93.5%) of white powder. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) δ 10.47(br s, 2H), 8.39 (br s, 1H), 7.40 (m, 7H), 5.33 (s, 2H), 2.26 (s, 3H).

#### **4-Bromo-3-(2-hydroxy-ethyl)-7-methyl-1H-indole-6-carboxylic acid benzyl ester**

A 1000 mL RBF was charged with 5-bromo-3-hydrazino-2-methyl benzoic acid benzyl ester hydrochloride (30.0 g, 0.081 mol), ethylene glycol (350 mL), and H<sub>2</sub>O (60 mL). This mixture was then heated to 40°C, and dihydrofuran (7.63 mL, 0.101 mol) was added. The reaction mixture was then heated to 100-105°C with stirring for 2.25 hours. The reaction was cooled to ambient temperature, diluted with EtOAc, washed with saturated NaCl (2x), dried (MgSO<sub>4</sub>), and concentrated. The resulting red oil was then chromatographed on SiO<sub>2</sub> eluting with a 10% EtOAc/DCM to 50% EtOAc/DCM gradient. Isolation of the product as an impure solid followed by trituration with DCM and hexanes afforded 8.45g (27.0%) of pale yellow solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.28 (br s, 1H), 7.94 (s, 1H), 7.40 (m, 6H), 5.36 (s, 1H), 3.96 (q, J = 6.4Hz, 2H), 3.28 (t, J = 6.4Hz, 2H), 2.72 (s, 3H),

#### **5-Bromo-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4,-b]indole-1,7-dicarboxylic acid 7-benzyl ester 1-ethyl ester**

To a solution of 4-bromo-3-(2-hydroxy-ethyl)-7-methyl-1*H*-indole-6-carboxylic acid benzyl ester (0.96 g, 2.5 mmol) in dichloroethane (60 mL) was added 2-oxo-pentanoic acid ethyl ester (0.57 g, 3.9 mmol) in dichloroethane (5 mL). The solution was heated at 70 °C and BF<sub>3</sub>-OEt<sub>2</sub> (0.80 mL, 6.3 mmol) was then added. The solution was cooled to room temperature and allowed to stir for 2 hours. The solution was diluted with brine, and extracted with EtOAc (3x), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. Purification of the residue by flash chromatography (SiO<sub>2</sub>, 2/1 hexanes/EtOAc) gave pyranoindole (781 mg, 61%) as an orange oil. APCI (M+H) 461 m/z. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.60 (s, 1H), 7.89 (s, 1H), 7.46-7.32 (m, 5H), 5.35 (s, 2H), 4.34-4.21 (m, 4H), 4.08-3.91 (m, 1H), 3.18-3.12 (m, 2H), 2.70 (s, 3H), 2.10-2.00 (m, 2H), 1.37-1.32 (m, 4H), 0.95 (t, *J* = 7.8 Hz, 3H).

**5-Cyano-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4,-*b*]indole-1,7-dicarboxylic acid 7-benzyl ester 1-ethyl ester**

A solution of 5-bromo-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4,-*b*]indole-1,7-dicarboxylic acid 7-benzyl ester 1-ethyl ester (861 mg, 1.7 mmol) in NMP (10 mL) was treated with CuCN (1.5 g, 17 mmol) at room temperature. The solution was then heated at 180 °C for 2 hours. The reaction was cooled to room temperature, diluted with H<sub>2</sub>O and EtOAc, and filtered through Celite. The filter pad was thoroughly rinsed with EtOAc. The filtrate was extracted with brine (2x), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. Purification of the residual oil by flash chromatography (SiO<sub>2</sub>, 3/1 hexanes/EtOAc) gave benzonitrile as a light yellow solid (680 mg, 87%). APCI (M+H) 461 m/z. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.69 (s, 1H), 8.12 (s, 1H), 7.48-7.36 (m, 5H), 5.37 (s, 2H), 4.33-4.26 (m, 3H), 3.99-3.91 (m, 1H), 3.13-3.02 (m, 2H), 2.83 (s, 3H), 2.20-2.14 (m, 1H), 2.12-2.02 (m, 1H), 1.43-1.33 (m, 1H), 1.34 (t, *J* = 7.2 Hz, 3H), 1.31-1.22 (m, 1H), 0.90 (t, *J* = 6.9 Hz, 3H).

**5-Cyano-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4,-*b*]indole-1,7-dicarboxylic acid 1-ethyl ester**

A solution of 5-cyano-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4,-*b*]indole-1,7-dicarboxylic acid 7-benzyl ester 1-ethyl ester (718 mg, 1.6 mmol) in a mixture of EtOAc/MeOH (30 mL/ 2/1) was treated with Pd(OH)<sub>2</sub> (280 mg) and subjected to H<sub>2</sub> using Parr shaker. After 16 hours, the solution was filtered through Celite. The filter cake was thoroughly rinsed with EtOAc and MeOH. The filtrate was concentrated to give the acid as a light yellow solid (551 mg, 93%). LCMS (M-H) 369 m/z. <sup>1</sup>H NMR (DMSO) δ 11.58 (s, 1H), 7.93 (s, 1H), 4.19-4.01 (m, 3H), 3.87-3.78 (m, 1H), 2.97-2.93 (m, 2H), 2.85 (s, 3H), 2.49-2.16 (m, 2H), 1.25-1.07 (m, 5H), 0.83 (t, *J* = 7.5 Hz, 3H).

**5-Cyano-7-hydroxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4,-*b*]indole-1-carboxylic acid ethyl ester**

A solution of 5-cyano-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4,-*b*]indole-1,7-dicarboxylic acid 1-ethyl ester (551 mg, 1.5 mmol) in THF (20 mL) was treated with BH<sub>3</sub>-THF (1M, 4.5 mL) at 0 °C. The solution was slowly warmed to room temperature. After 2 hours, the reaction mixture was diluted with 1M HCl and extracted with EtOAc (3x). The combined organics were washed with saturated NaHCO<sub>3</sub>, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. Purification of the residue by flash chromatography (SiO<sub>2</sub>, 1/1 hexanes/EtOAc) provided 350 mg (66 %) of the alcohol as a light yellow foam. GCMS M<sup>+</sup> 356 m/z. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.77 (s, 1H), 7.77 (s, 1H), 4.73 (s, 2H), 4.34-4.24 (m, 3H), 3.99-3.90 (m, 1H), 3.17-3.03 (m, 2H), 2.38 (s, 3H), 2.35 (bs, 1H), 2.21-1.99 (m, 2H), 1.44-1.23 (m, 5H), 0.89 (t, *J* = 7.8 Hz, 3H).

**5-Cyano-7-formyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4,-*b*]indole-1-carboxylic acid ethyl ester**

A solution of 5-cyano-7-hydroxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4,-*b*]indole-1-carboxylic acid ethyl ester (350 mg, 0.98 mmol) in dichloroethane (20 mL) was treated with activated MnO<sub>2</sub> (0.58 g, 6.7 mmol) at room temperature. The solution was then heated at reflux for 1 hour. The mixture was filtered through a Celite/silica gel plug which was thoroughly rinsed with EtOAc. The filtrate was concentrated to give 341 mg (98%) of the aldehyde as a light yellow solid. GCMS M<sup>+</sup> 354 m/z. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 10.28 (s, 1H), 8.92 (s,

1H), 7.92 (s, 1H), 4.35-4.24 (m, 3H), 3.98-3.92 (m, 1H), 3.16-3.09 (m, 2H), 2.87 (s, 3H), 2.39-2.00 (m, 2H), 1.45-1.37 (m, 4H), 0.91 (t,  $J = 7.5$  Hz, 3H).

**5-Cyano-7-hydroxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indole-1-carboxylic acid ethyl ester**

A solution of 5-cyano-7-formyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indole-1-carboxylic acid ethyl ester (146 mg, 0.36 mmol) in *t*-butanol (15 mL) was treated with  $\text{SeO}_2$  (13 mg, 0.11 mmol) followed by  $\text{H}_2\text{O}_2$  (30%, 0.2 mL) and TFA (2 drops). After 1.5 hours, additional  $\text{SeO}_2$  (12 mg) and  $\text{H}_2\text{O}_2$  (0.2 mL) were added. After 4 hours, the solution was diluted with buffer (pH 6.9) and extracted with EtOAc. The combined organics were dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated. The crude material was used without further purification. LCMS (M-H) 423 m/z. The residue was diluted with MeOH (10 mL) and treated with 10%  $\text{K}_2\text{CO}_3$  at room temperature. After 1 hour, the solution was concentrated. The material was extracted with EtOAc and 1M HCl (2x), dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to provide an off-white solid (47 mg, 33%-2 steps). LCMS (M-H) 341 m/z.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  8.20 (s, 1H), 6.97 (s, 1H), 5.10 (s, 1H), 4.30-4.23 (m, 3H), 3.99-3.91 (m, 1H), 3.13-2.99 (m, 2H), 2.40 (s, 3H), 2.17-1.93 (m, 2H), 1.42-1.29 (m, 5H), 0.89 (t,  $J = 7.8$  Hz, 3H).

**5-Cyano-7-(2-methoxy-ethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indole-1-carboxylic acid**

A solution of 5-cyano-7-hydroxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indole-1-carboxylic acid ethyl ester (50 mg, 0.15 mmol) in DMF (5 mL) was added to a solution of bromoethyl-methylether (0.1 mL, 1.0 mmol) in DMF (2 mL) containing  $\text{Cs}_2\text{CO}_3$  (20 mg, 61 mmol) at room temperature. The reaction was heated at 60 °C for 1 hour, cooled to room temperature, diluted with brine, and extracted with EtOAc (3x). The combined organics were dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated. Purification of the residue by flash chromatography ( $\text{SiO}_2$ , 2/1-1/1 hexanes/ EtOAc) provided the ether as a yellow oil that was used without further purification. LCMS (M-H) 399.

A solution of the ethyl ester (30 mg, 0.07 mmol) in EtOH (5 mL) was treated with 1M NaOH (1 mL) at room temperature. After 16 hours, the solution was concentrated, diluted with H<sub>2</sub>O, and extracted with EtOAc (2x). The combined aqueous layers were made acidic with 1M HCl, extracted with EtOAc, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The residue was purified by flash chromatography (SiO<sub>2</sub>, 10% MeOH/DCM-3%TFA/10% MeOH/CHCl<sub>3</sub>) followed by trituration with Et<sub>2</sub>O/hexanes to give 7 mg (61%) of the formic acid as an off-white solid. LCMS (M-H) 371 m/z. <sup>1</sup>H NMR (CDCl<sub>3</sub>+CD<sub>3</sub>OD) δ 8.99 (s, 1H), 7.06 (s, 1H), 4.17-4.14 (m, 1H), 3.80-3.76 (m, 2H), 3.47 (s, 3H), 3.05 (m, 2H), 2.94 (m, 7H), 2.43 (s, 3H), 2.14-2.09 (m, 1H), 1.37-1.35 (m, 2H), 0.89 (t, *J* = 7.2 Hz, 3H).

### Example 32

#### 5-Cyano-8-methyl-7-(5-methyl-isoxazol-3-ylmethoxy)-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indole-1-carboxylic acid

A solution of 5-cyano-7-hydroxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indole-1-carboxylic acid ethyl ester (25 mg, 0.07 mmol) in DMF (5 mL) was treated with Cs<sub>2</sub>CO<sub>3</sub> (52 mg, 0.16 mmol) at room temperature. The solution was treated with a mixture of bromomethyl isoxazole (20 mg, 0.11 mmol) in DMF (2 mL) and then heated at 60 °C for 1.5 hours. The solution was cooled to room temperature and diluted with brine, extracted with EtOAc (3x), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. Purification of the residue by flash chromatography (SiO<sub>2</sub>, 2/1 hexanes/EtOAc) provided the ether as a yellow oil that was used without further purification. LCMS (M-H) 436 m/z.

A solution of the ethyl ester (36 mg, 0.08 mmol) in EtOH (10 mL) was treated with 1M NaOH (1 mL) at room temperature. The solution was heated at 60 °C for 0.5 hours and then concentrated. The residue was diluted with H<sub>2</sub>O, acidified with 1M HCl, and extracted with EtOAc (3x). The combined organics were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The material was purified by preparative LCMS which provided the formic acid as an off-white fluffy solid (12 mg, 36%) following lyophilization. LCMS (M-H) 408 m/z. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.38 (bs, 1H), 7.13 (s, 1H), 6.10 (s, 1H), 5.14 (s, 2H), 4.22-4.11 (m, 2H), 3.16-3.12 (m, 2H), 2.45

(s, 3H), 2.40 (s, 3H), 2.25-2.15 (m, 1H), 2.11-2.00 (m, 1H), 1.42-1.38 (m, 2H), 0.94 (t,  $J = 7.8$  Hz, 3H).

### Example 33

**(1*R*\*,10*S*)- [1-sec-Butyl-5-cyano-7-(2-ethoxy-ethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid**

**(1*R*\*,10*S*)-5-Bromo-1-sec-butyl-1-ethoxycarbonylmethyl-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indole-7-carboxylic acid benzyl ester**

To a solution of 4-bromo-3-(2-hydroxy-ethyl)-7-methyl-1*H*-indole-6-carboxylic acid benzyl ester (2.00 g, 5.15 mmol) and (*S*)-ethyl 4-methyl-3-oxohexanoate (0.89 g, 5.15 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (50 mL) was added BF<sub>3</sub>·OEt<sub>2</sub> (0.80 g, 5.66 mmol) dropwise at room temperature. The solution was stirred for 4 hours then quenched by the addition of saturated aqueous Na<sub>2</sub>CO<sub>3</sub> (10 mL). The reaction mixture was diluted with water and extracted three times with diethyl ether (250 mL). The combined organic layers were washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to dryness. The crude product was purified by silica gel chromatography (5% to 20% EtOAc/hexane) to afford 0.77 g (28%) of the esters as an ~1:1 mixture of 2 diastereomers. <sup>1</sup>H NMR (CDCl<sub>3</sub>): ~1:1 mixture of 2 diastereomers,  $\delta$  9.78 (br s, 0.5H, indole NH of diastereomer 1), 9.76 (br s, 0.5H, indole NH of diastereomer 2), 7.88 (s, 1H), 7.31-7.47 (m, 5H), 5.35 (s, 2H), 4.27-4.05 (m, 3H), 3.73 (m, 1H), 3.21-2.96 (m, 4.5H), 2.72 (s, 3H), 2.16 (m, 1H), 1.66 (m, 0.5H), 1.35 (m, 0.5H), 1.28 (t,  $J = 7$  Hz, 3H), 1.10 (m, 0.5H), 1.03 (d,  $J = 7$  Hz, 1.5H), 0.93 (t,  $J = 7$  Hz, 1.5H), 0.73 (t,  $J = 7$  Hz, 1.5H), 0.63 (d,  $J = 7$  Hz, 1.5H).

**(1*R*\*,10*S*)-1-sec-Butyl-5-cyano-1-ethoxycarbonylmethyl-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indole-7-carboxylic acid benzyl ester**

A mixture of (1*R*\*,10*S*)-5-bromo-1-sec-butyl-1-ethoxycarbonylmethyl-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indole-7-carboxylic acid benzyl ester (0.77 g, 1.41 mmol) and CuCN (0.63 g, 7.09 mmol) in *N*-methyl-2-pyrrolidinone (30 mL) was immersed in a pre-heated oil bath



(190 °C). After 2 hours, the reaction was complete by TLC analysis and cooled to ambient temperature. The reaction mixture was diluted with water (30 mL) and diethyl ether (30 mL) and filtered through a pad of Celite. The filtrate was extracted three times with diethyl ether (150 mL). The combined organic layers were washed with water (5x) and brine (1x), dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to afford the crude ester as an ~1:1 mixture of 2 diastereomers which was used directly in the next reaction. <sup>1</sup>H NMR (CDCl<sub>3</sub>): ~1:1 mixture of 2 diastereomers, δ 10.05 (br s, 0.5H, indole NH of diastereomer 1), 10.03 (br s, 0.5H, indole NH of diastereomer 2), 8.11 (s, 1H), 7.48-7.35 (m, 5H), 5.36 (s, 2H), 4.28-4.09 (m, 3H), 3.37 (m, 0.5H), 3.17-2.99 (m, 4H), 2.83 (s, 3H), 2.36 (m, 0.5H), 2.16 (m, 1H), 2.01 (m, 0.5H), 1.66 (m, 0.5H), 1.36 (m, 0.5H), 1.26 (t, J = 7 Hz, 3H), 1.14 (m, 0.5H), 1.05 (d, J = 6 Hz, 1.5H), 0.93 (t, J = 8 Hz, 1.5H), 0.76 (t, J = 7 Hz, 1.5H), 0.65 (d, J = 6 Hz, 1.5H).

**(1*R*\*,10*S*)-1-sec-Butyl-5-cyano-1-ethoxycarbonylmethyl-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indole-7-carboxylic acid**

A solution containing crude (1*R*\*,10*S*)-1-sec-butyl-5-cyano-1-ethoxycarbonylmethyl-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indole-7-carboxylic acid benzyl ester (0.81 g) and 10% Pd –on Carbon (0.2 g) in EtOAc (20 mL) was stirred under a balloon of hydrogen gas for 3 hours at ambient temperature. The reaction mixture was filtered through a pad of Celite®, washed with methanol and concentrated to afford the crude acid as an ~1:1 mixture of 2 diastereomers which was used directly in the next reaction. <sup>1</sup>H NMR (CDCl<sub>3</sub>): ~1:1 mixture of 2 diastereomers, δ 10.12 (br s, 0.5H, indole NH of diastereomer 1), 10.10 (br s, 0.5H, indole NH of diastereomer 2), 8.20 (s, 1H), 4.30-4.08 (m, 2.5H), 3.78 (m, 1H), 3.40 (t, J = 7 Hz, 1H), 3.19-2.98 (m, 3.5H), 2.88 (s, 3H), 2.43 (app t, J = 7 Hz, 0.5H), 2.20 (m, 0.5H), 2.03 (m, 0.5H), 1.68 (m, 0.5H), 1.36 (m, 0.5H), 1.29 (t, J = 7 Hz, 3H), 1.14 (m, 0.5H), 1.06 (d, J = 6 Hz, 1.5H), 0.94 (t, J = 8 Hz, 1.5H), 0.75 (t, J = 7 Hz, 1.5H), 0.65 (d, J = 6 Hz, 1.5H).

**(1*R*\*,10*S*)-(1-sec-Butyl-5-cyano-7-hydroxymethyl-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl)-acetic acid ethyl ester**

To a solution of crude (1*R*\*,10*S*)-1-sec-butyl-5-cyano-1-ethoxycarbonylmethyl-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indole-7-carboxylic acid (0.6 g) in THF (20 mL) at 0 °C was added BH<sub>3</sub>·THF (1.0M in THF, 3 mL, 3.00 mmol). The cooling bath was removed, and the solution was stirred for 1 hour. The reaction mixture was re-cooled to 0 °C and a second portion of BH<sub>3</sub>·THF (1.0M in THF, 3 mL, 3.00 mmol) was added. The reaction mixture was stirred an additional 1 hour at ambient temperature then carefully quenched with water. The reaction mixture was diluted with 1.0 N HCl and EtOAc, and the layers were separated. The organic layer was washed with water and brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to afford the crude alcohol as an ~1:1 mixture of 2 diastereomers which was used directly in the next reaction. <sup>1</sup>H NMR (CDCl<sub>3</sub>): ~1:1 mixture of 2 diastereomers, δ 9.66 (br s, 1H, indole NH), 7.43 (s, 1H), 4.80 (s, 2H), 4.27-4.06 (m, 3H), 3.77 (m, 1H), 3.16-2.98 (m, 3H), 2.57 (s, 3H), 2.19 (m, 0.5H), 1.63 (m, 0.5H), 1.34 (m, 0.5H), 1.28 (t, J = 7 Hz, 3H), 1.18 (m, 0.5H), 1.04 (d, J = 6 Hz, 1.5H), 0.93 (t, J = 8 Hz, 1.5H), 0.74 (t, J = 7 Hz, 1.5H), 0.63 (d, J = 6 Hz, 1.5H).

**(1*R*\*,10*S*)-(1-sec-Butyl-5-cyano-7-formyl-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl)-acetic acid ethyl ester**

To a solution of crude (1*R*\*,10*S*)-(1-sec-butyl-5-cyano-7-hydroxymethyl-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl)-acetic acid ethyl ester (0.57 g) in anhydrous benzene (20 mL) was added a homogeneous mixture of activated 85% MnO<sub>2</sub> (<5 μ particle size, 0.3 g, 2.96 mmol) and Celite® (0.5 g). The mixture was immersed in a pre-heated oil bath (60 °C) and agitated vigorously. After 1 hour, a second portion of MnO<sub>2</sub> (1.0 g) and Celite® (1.0 g) was added and heating was continued for 1 hour. The reaction mixture was cooled to ambient temperature then filtered through Celite®. The solids were washed with EtOAc, and the yellow filtrate was concentrated to afford the crude aldehyde as an ~1:1 mixture of 2 diastereomers which was used directly in the next reaction. <sup>1</sup>H NMR (CDCl<sub>3</sub>): ~1:1 mixture of 2 diastereomers, δ 10.27 (s, 1H), 7.90 (s, 1H), 4.30-4.11 (m, 3H), 3.79 (m, 1H), 3.18-3.02 (m, 3H), 2.88 (s, 3H), 2.17 (m, 0.5H), 1.67 (m, 0.5H), 1.36 (m, 0.5H), 1.30 (t, J = 7 Hz, 3H), 1.23 (m, 0.5H), 1.06 (d, J = 6 Hz, 1.5H), 0.94 (t, J = 8 Hz, 1.5H), 0.75 (t, J = 7 Hz, 1.5H), 0.64 (d, J = 6 Hz, 1.5H).

**(1*R*\*,10*S*)-(1-sec-Butyl-5-cyano-7-hydroxy-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl)-acetic acid ethyl ester**

To a solution of (1*R*\*,10*S*)-(1-sec-butyl-5-cyano-7-formyl-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl)-acetic acid ethyl ester (0.53 g) in *tert*-butanol (10 mL) was added SeO<sub>2</sub> (46 mg, 0.41 mmol) followed by 30% H<sub>2</sub>O<sub>2</sub> (5 mL) and trifluoroacetic acid (20 μL). The reaction mixture was stirred at ambient temperature for 1 hour, and then diluted with EtOAc, washed with saturated NaHCO<sub>3</sub> (2x), water and brine; dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>; filtered and concentrated to afford the crude formate ester as an ~1:1 mixture of 2 diastereomers which was used directly in the next reaction.

The crude residue was dissolved in MeOH (15 mL) and to this was added 10% aqueous K<sub>2</sub>CO<sub>3</sub> (10 mL). After 1 hour, the reaction mixture was diluted with EtOAc, washed with water (2x) and brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The crude phenol was purified by silica gel chromatography (20% to 30% EtOAc/hexane) to afford 0.19 g of (1*R*\*,10*S*)-(1-sec-butyl-5-cyano-7-hydroxy-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl)-acetic acid ethyl ester (36% over 6 steps) as an ~1:1 mixture of diastereomers. <sup>1</sup>H NMR (CDCl<sub>3</sub>): ~1:1 mixture of 2 diastereomers, δ 9.38 (br s, 0.5H, indole NH of diastereomer 1), 9.36 (br s, 0.5H, indole NH of diastereomer 2), 6.96 (s, 1H), 5.26 (br s, 1H), 4.26-4.06 (m, 3H), 3.76 (m, 1H), 3.15-2.96 (m, 3H), 2.41 (s, 3H), 2.16 (m, 0.5H), 1.68 (m, 0.5H), 1.35 (m, 0.5H), 1.27 (t, *J* = 7 Hz, 3H), 1.14 (m, 0.5H), 1.03 (d, *J* = 6 Hz, 1.5H), 0.93 (t, *J* = 8 Hz, 1.5H), 0.75 (t, *J* = 7 Hz, 1.5H), 0.65 (d, *J* = 6 Hz, 1.5H).

**(1*R*\*,10*S*)-[1-sec-Butyl-5-cyano-7-(2-ethoxy-ethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid ethyl ester**

To a solution containing (1*R*\*,10*S*)-(1-sec-butyl-5-cyano-7-hydroxy-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl)-acetic acid ethyl ester (50 mg, 0.13 mmol) in anhydrous DMF (5 mL) was added Cs<sub>2</sub>CO<sub>3</sub> (66 mg, 0.20 mmol) and 1-bromo-2-ethoxyethane (23 μL, 0.20 mmol). The reaction mixture was immersed in a pre-heated oil bath (60 °C). After 2 hours, additional 1-bromo-2-ethoxyethane (23 μL, 0.20 mmol) was added, and the reaction mixture was

maintained at 60 °C for 3 hours. The reaction mixture was cooled to ambient temperature and diluted with diethyl ether and water. The layers were separated, and the organic phase was washed with water (5x) and brine (1x), dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to afford the crude product as an ~1:1 mixture of 2 diastereomers which was used directly in the next reaction. <sup>1</sup>H NMR (CDCl<sub>3</sub>): ~1:1 mixture of 2 diastereomers, δ 9.40 (br s, 0.5H, indole NH of diastereomer 1), 9.38 (br s, 0.5H, indole NH of diastereomer 2), 7.06 (s, 1H), 4.26-4.01 (m, 5H), 3.82-3.72 (m, 3H), 3.62 (t, J = 7 Hz, 2H), 3.15-2.97 (m, 3H), 2.42 (s, 3H), 2.17 (m, 0.5H), 1.66 (m, 0.5H), 1.35 (m, 0.5H), 1.32-1.18 (m, 6H), 1.14 (m, 0.5H), 1.03 (d, J = 6 Hz, 1.5H), 0.93 (t, J = 8 Hz, 1.5H), 0.75 (t, J = 7 Hz, 1.5H), 0.65 (d, J = 6 Hz, 1.5H) ppm.

**(1*R*\*,10*S*)-[1-sec-Butyl-5-cyano-7-(2-ethoxy-ethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid**

To a solution containing crude (1*R*\*,10*S*)-[1-sec-butyl-5-cyano-7-(2-ethoxy-ethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid ethyl ester (~50 mg) in ethanol (4 mL) and THF (1 mL) was added 3 N NaOH (1 mL) at ambient temperature. After 5 hours, the reaction mixture was diluted with water and washed with diethyl ether. The aqueous phase was acidified with 1 N HCl, and the product was extracted with diethyl ether. The ether extract was washed with 1 N HCl and brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The crude product was purified by silica gel chromatography (1% to 20% MeOH/DCM) and the purified acid was triturated with diethyl ether/hexane. The product was collected on a Hirsch funnel, washed with hexane then dried *in vacuo* to afford 34 mg of (1*R*\*,10*S*)-[1-sec-butyl-5-cyano-7-(2-ethoxy-ethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid (62% over 2 steps) as an ~1:1 mixture of 2 diastereomers. ESI MS *m/z* 413.2 (M-H).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): ~1:1 mixture of 2 diastereomers, δ 9.13 (br s, 0.5H, indole NH of diastereomer 1), 9.09 (br s, 0.5H, indole NH of diastereomer 2), 7.05 (s, 1H), 4.18-4.00 (m, 2H), 3.97-3.77 (m, 3H), 3.63 (t, J = 7 Hz, 2H), 3.50 (t, J = 7 Hz, 2H), 3.13-3.03 (m, 4H), 2.27 (s, 3H), 2.17-1.61 (m, 2H), 1.36-1.06 (m, 5H), 1.04 (d, J = 6 Hz, 1.5H), 0.93 (t, J = 8 Hz, 1.5H), 0.75 (t, J = 7 Hz, 1.5H), 0.65 (d, J = 6 Hz, 1.5H).

**Example 34**

**(1*R*\*,1*S*)- [1-sec-Butyl-5-cyano-7-(2-isopropoxy-ethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid****(1*R*\*,1*S*)- [1-sec-Butyl-5-cyano-7-(2-isopropoxy-ethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid ethyl ester**

To a solution containing (1*R*\*,1*S*)-(1-sec-butyl-5-cyano-7-hydroxy-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl)-acetic acid ethyl ester (50 mg, 0.13 mmol) in anhydrous DMF (5 mL) was added Cs<sub>2</sub>CO<sub>3</sub> (66 mg, 0.20 mmol) and 1-bromo-2-isopropoxyethane (~40  $\mu$ L, >0.20 mmol). The reaction mixture was immersed in a pre-heated oil bath (60 °C). After 4 hours, additional 1-bromo-2-isopropoxyethane (~40  $\mu$ L, >0.20 mmol) was added, and the reaction mixture was maintained at 60°C for 3 hours. The reaction mixture was cooled to ambient temperature and diluted with diethyl ether and water. The layers were separated, and the organic phase was washed with water (5x) and brine (1x), dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The crude ester was purified by silica gel chromatography (25% EtOAc/hexane) to afford the ester as an ~1:1 mixture of 2 diastereomers. <sup>1</sup>H NMR (CDCl<sub>3</sub>): ~1:1 mixture of 2 diastereomers,  $\delta$  9.40 (br s, 0.5H, indole NH of diastereomer 1), 9.38 (br s, 0.5H, indole NH of diastereomer 2), 7.06 (s, 1H), 4.35-3.92 (m, 5H), 3.79-3.57 (m, 3H), 3.44 (m, 2H), 3.19-2.80 (m, 3H), 2.40 (s, 3H), 2.17 (m, 0.5H), 1.66 (m, 0.5H), 1.39-1.09 (m, 12H), 1.03 (d, J = 6 Hz, 1.5H), 0.93 (t, J = 8 Hz, 1.5H), 0.75 (m, 3H).

**(1*R*\*,1*S*)- [1-sec-Butyl-5-cyano-7-(2-isopropoxy-ethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid**

To a solution containing crude (1*R*\*,1*S*)- [1-sec-butyl-5-cyano-7-(2-isopropoxy-ethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid ethyl ester (~50 mg) in ethanol (4 mL) and THF (1 mL) was added 3 N NaOH (1 mL) at ambient temperature. After 5 hours, the reaction mixture was diluted with water and washed with diethyl ether. The aqueous phase was acidified with 1 N HCl, and the product was extracted with diethyl ether. The ether extract was washed with 1 N HCl and brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The crude product was purified by silica gel chromatography (1% to 20%

MeOH/DCM), and the purified acid was triturated with diethyl ether/hexane. The product was collected on a Hirsch funnel, washed with hexane, and dried *in vacuo* to afford 16 mg of (1*R*\*,10*S*)- [1-sec-butyl-5-cyano-7-(2-isopropoxy-ethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid (29% over 2 steps) as an ~1:1 mixture of 2 diastereomers. ESI MS *m/z* 429.2 (M+H)<sup>+</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>): ~1:1 mixture of 2 diastereomers, δ 8.88 (br s, 0.5H, indole NH of diastereomer 1), 8.83 (br s, 0.5H, indole NH of diastereomer 2), 7.06 (s, 1H), 4.19-4.11 (m, 3H), 3.89 (m, 1H), 3.78 (app t, J = 5 Hz, 2H), 3.69 (m, 1H), 3.12-3.04 (m, 4H), 2.32 (s, 3H), 2.17 (m, 1H), 1.65 (m, 1H), 1.27 (m, 1H), 1.21 (d, J = 5 Hz, 6H), 1.05 (d, J = 7 Hz, 1.5H), 0.93 (t, J = 8 Hz, 1.5H), 0.77 (m, 3 H).

### Example 35

**(1*R*\*,10*S*)- [1-sec-Butyl-5-cyano-7-(5-dimethylamino-[1,2,4]thiadiazol-3-ylmethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid**

**(1*R*\*,10*S*)-[1-sec-Butyl-5-cyano-7-(5-dimethylamino-[1,2,4]thiadiazol-3-ylmethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid ethyl ester**

To a solution containing (1*R*\*,10*S*)-(1-sec-butyl-5-cyano-7-hydroxy-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl)-acetic acid ethyl ester (43 mg, 0.11 mmol) in anhydrous NMP (2 mL) was added Cs<sub>2</sub>CO<sub>3</sub> (57 mg, 0.17 mmol), KI (19 mg, cat.), and 3-chloromethyl-5-dimethylamino-1,2,4-thiadiazole (27 mg, 0.17 mmol). The reaction mixture was maintained at ambient temperature. After 16 hours, the reaction mixture was diluted with diethyl ether and 10% aqueous HCl. The layers were separated, and the organic phase was washed with water (5 x) and brine (1x), dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The crude ester was purified by silica gel chromatography (30% to 60% EtOAC/hexane) to afford 46 mg of the ester as an ~1:1 mixture of 2 diastereomers. <sup>1</sup>H NMR (CDCl<sub>3</sub>): ~1:1 mixture of 2 diastereomers, δ 9.42 (br s, 0.5H, indole NH of diastereomer 1), 9.39 (br s, 0.5H, indole NH of diastereomer 2), 7.18 (s, 1H), 5.11 (s, 2H), 4.30-4.04 (m, 3H), 3.75 (m, 1H), 3.39 (t, J = 7 Hz, 1H), 3.18 (s, 6H), 3.13-2.95 (m, 3H), 2.44 (s, 3H), 2.39 (m, 1H), 2.18 (m, 1H), 2.01 (m, 1H), 1.66 (m, 0.5H), 1.30 (t, J = 7 Hz, 3H), 1.35 (m, 0.5H), 1.16 (m, 0.5H), 1.03 (d, J = 6 Hz, 1.5H), 0.93 (t, J = 8 Hz, 1.5H), 0.75 (t, J = 7 Hz, 1.5H), 0.63 (d, J = 7 Hz, 1.5H).

**(1*R*\*,10*S*)-[1-sec-Butyl-5-cyano-7-(5-dimethylamino-[1,2,4]thiadiazol-3-ylmethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid**

To a solution containing (1*R*\*,10*S*)-[1-sec-butyl-5-cyano-7-(5-dimethylamino-[1,2,4]thiadiazol-3-ylmethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid ethyl ester (46 mg) in ethanol (2 mL) and THF (1 mL) was added 3 N NaOH (1 mL) at ambient temperature. After 16 hours, the reaction mixture was diluted with water and acidified with 1 N HCl. The product was extracted with diethyl ether. The ether extract was washed with 1 N HCl and brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The crude product was triturated with diethyl ether/hexane. The product was collected on a Hirsch funnel, washed with hexane, and dried *in vacuo* to afford 29 mg of (1*R*\*,10*S*)-[1-sec-butyl-5-cyano-7-(5-dimethylamino-[1,2,4]thiadiazol-3-ylmethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid (54% over 2 steps) as an ~1:1 mixture of 2 diastereomers. ESI MS *m/z* 484.1 (M+H)<sup>+</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>): ~1:1 mixture of 2 diastereomers, δ 9.04 (br s, 0.5H, indole NH of diastereomer 1), 8.98 (br s, 0.5H, indole NH of diastereomer 2), 7.15 (s, 1H), 5.09 (s, 2H), 4.12 (m, 1H), 3.88 (m, 1H), 3.19 (s, 6H), 3.13-2.98 (m, 3H), 2.38 (s, 3H), 2.20 (m, 1H), 1.68 (m, 1H), 1.28 (m, 1H), 1.12 (m, 1H), 1.09 (d, *J* = 6 Hz, 1.5H), 0.93 (t, *J* = 8 Hz, 1.5H), 0.75 (t, *J* = 7 Hz, 1.5H), 0.63 (d, *J* = 7 Hz, 1.5H).

**Example 36**

**(1*R*\*,10*S*)-[1-sec-Butyl-5-cyano-7-(1,5-dimethyl-1*H*-pyrazol-3-ylmethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid**

**(1*R*\*,10*S*)-[1-sec-Butyl-5-cyano-7-(1,5-dimethyl-1*H*-pyrazol-3-ylmethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid ethyl ester**

To a solution containing (1*R*\*,10*S*)-(1-sec-butyl-5-cyano-7-hydroxy-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl)-acetic acid ethyl ester (47 mg, 0.12 mmol) in anhydrous NMP (2 mL) was added Cs<sub>2</sub>CO<sub>3</sub> (83 mg, 0.25 mmol) and 3-bromomethyl-1,5-dimethylpyrazole (48 mg, 0.25 mmol). The reaction mixture was maintained at ambient temperature. After 16 hours, the reaction mixture was diluted with diethyl ether and 10% aqueous HCl. The layers

were separated, and the organic phase was washed with water (5x) and brine (1x), dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The crude ester was purified by silica gel chromatography (50% to 60% EtOAc/hexane) to afford the ester as an ~1:1 mixture of 2 diastereomers. <sup>1</sup>H NMR (CDCl<sub>3</sub>): ~1:1 mixture of 2 diastereomers, δ 9.37 (br s, 0.5H, indole NH of diastereomer 1), 9.35 (br s, 0.5H, indole NH of diastereomer 2), 7.16 (s, 1H), 6.10 (s, 1H), 5.02 (s, 2H), 4.26-4.06 (m, 3H), 3.77 (s, 3H), 3.74 (m, 2H), 3.15-2.96 (m, 3H), 2.41 (s, 3H), 2.27 (s, 3H), 2.17 (m, 1H), 1.66 (m, 0.5H), 1.35 (m, 0.5H), 1.26 (t, J = 7 Hz, 3H), 1.16 (m, 0.5H), 1.03 (d, J = 6 Hz, 1.5H), 0.94 (t, J = 8 Hz, 1.5H), 0.76 (t, J = 7 Hz, 1.5H), 0.65 (d, J = 7 Hz, 1.5H).

**(1*R*\*,10*S*)-[1-sec-Butyl-5-cyano-7-(1,5-dimethyl-1*H*-pyrazol-3-ylmethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid**


To a solution containing (1*R*\*,10*S*)-[1-sec-butyl-5-cyano-7-(1,5-dimethyl-1*H*-pyrazol-3-ylmethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid ethyl ester (~50 mg) in ethanol (4 mL) and THF (1 mL) was added 3 N NaOH (2 mL) at ambient temperature. After 16 hours, the reaction mixture was diluted with water and acidified with 1 N HCl. The product was extracted with diethyl ether. The ether extract was washed with 1 N HCl and brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The crude product was triturated with diethyl ether/hexane. The product was collected on a Hirsch funnel, washed with hexane then dried *in vacuo* to afford 27 mg of (1*R*\*,10*S*)-[1-sec-butyl-5-cyano-7-(1,5-dimethyl-1*H*-pyrazol-3-ylmethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid (50% over 2 steps) as an ~1:1 mixture of 2 diastereomers. ESI MS *m/z* 449.2 (M-H)<sup>-</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>): ~1:1 mixture of 2 diastereomers, δ 9.16 (br s, 0.5H, indole NH of diastereomer 1), 9.13 (br s, 0.5H, indole NH of diastereomer 2), 7.06 (s, 1H), 6.12 (s, 1H), 4.99 (m, 2H), 4.13-4.08 (m, 1H), 3.85 (m, 1H), 3.78 (s, 3H), 3.12 (m, 2H), 3.00 (m, 3H), 2.28 (s, 3H), 2.23 (s, 3H), 1.70 (m, 0.5H), 1.35 (m, 0.5H), 1.16 (m, 0.5H), 1.06 (d, J = 6 Hz, 1.5H), 0.94 (t, J = 8 Hz, 1.5H), 0.75 (t, J = 7 Hz, 1.5H), 0.68 (d, J = 7 Hz, 1.5H).

By appropriate selection of suitable starting materials, other compounds of the invention may be prepared according to the procedures described in the foregoing examples.

















Representative examples of further pyranoindole derivatives and analogues thus prepared are set forth in Table 2 below.










Table 2. Pyranoindole derivatives





















R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, R<sub>8</sub> = H if unspecified










Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>7</sub>	R <sub>9</sub>	R <sub>10</sub>	Y	M	Mass Spec* (m/z)
1	H	n-Propyl	CN	H	CH <sub>3</sub>	CH <sub>2</sub>	CH <sub>2</sub>	341 (M-H) <sup>+</sup>
2	H	n-Propyl	CN	CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>2</sub>	CH <sub>2</sub>	355 (M-H) <sup>+</sup>
3	H	n-Propyl	CN	CH <sub>2</sub> CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>2</sub>	CH <sub>2</sub>	369 (M-H) <sup>+</sup>
4	H	n-Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	CH <sub>2</sub>	383 (M-H) <sup>+</sup>
5	H	n-Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	CH <sub>2</sub>	-
6	H	n-Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	CH <sub>2</sub>	395 (M-H) <sup>+</sup>
7	H	n-Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	CH <sub>2</sub>	423 (M-H) <sup>+</sup>










Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>7</sub>	R <sub>9</sub>	R <sub>10</sub>	Y	M	Mass Spec* (m/z)
8	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	CH <sub>2</sub>	395 (M-H) <sup>+</sup>
9	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	CH <sub>2</sub>	409 (M-H) <sup>+</sup>
10	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	CH <sub>2</sub>	423 (M-H) <sup>+</sup>
11	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	CH <sub>2</sub>	393 (M-H) <sup>+</sup>
12	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	CH <sub>2</sub>	425 (M-H) <sup>+</sup>
13	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	CH <sub>2</sub>	411 (M-H) <sup>+</sup>
14	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	CH <sub>2</sub>	411 (M-H) <sup>+</sup>
15	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	CH <sub>2</sub>	417 (M-H) <sup>+</sup>
16	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	CH <sub>2</sub>	475 (M-H) <sup>+</sup>
17	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	CH <sub>2</sub>	459 (M-H) <sup>+</sup>

Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>7</sub>	R <sub>9</sub>	R <sub>10</sub>	Y	M	Mass Spec* (m/z)
18	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	CH <sub>2</sub>	437 (M-H) <sup>-</sup>
19	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	CH <sub>2</sub>	466 (M-H) <sup>-</sup>
20	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	CH <sub>2</sub>	431 (M-H) <sup>-</sup>
21	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	CH <sub>2</sub>	435 (M-H) <sup>-</sup>
22	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	383 (M+H) <sup>+</sup>
23	H	<i>n</i> -Propyl (1 <i>R</i> )	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	424 (M+H) <sup>+</sup>
24	H	<i>n</i> -Propyl (1 <i>R</i> )	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	-
25	H	<i>n</i> -Propyl (1 <i>R</i> )	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	435 (M-H) <sup>-</sup>
26	H	<i>n</i> -Propyl (1 <i>R</i> )	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	413 (M-H) <sup>-</sup>









Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>7</sub>	R <sub>9</sub>	R <sub>10</sub>	Y	M	Mass Spec* (m/z)
27	H	<i>n</i> -Propyl (1 <i>R</i> )	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	399 (M-H) <sup>-</sup>
28	H	<i>n</i> -Propyl (1 <i>R</i> )	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	399 (M-H) <sup>-</sup>
29	H	<i>n</i> -Propyl (1 <i>R</i> )	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	441 (M+H) <sup>+</sup>
30	H	<i>n</i> -Propyl (1 <i>R</i> )	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	470 (M+H) <sup>+</sup>
31	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	bond	bond	371 (M-H) <sup>-</sup>
32	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	bond	bond	408 (M-H) <sup>-</sup>
33	H	SecButyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	413 (M-H) <sup>-</sup>
34	H	SecButyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	429 (M+H) <sup>+</sup>
35	H	SecButyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	484 (M-H) <sup>-</sup>








Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>7</sub>	R <sub>9</sub>	R <sub>10</sub>	Y	M	Mass Spec* (m/z)
36	H	SecButyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	449 (M-H) <sup>-</sup>
37	H	n-Propyl	CN		CH <sub>3</sub>	bond	bond	404 (M-H) <sup>-</sup>
38	H	n-Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	420 (M+H) <sup>+</sup>
39	H	n-Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	399 (M+H) <sup>+</sup>
40	H	n-Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	-
41	H	n-Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	420 (MH) <sup>+</sup>
42	H	n-Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	398 (M+H) <sup>+</sup>
43	H	n-Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	387 (MH) <sup>+</sup>
44	H	n-Propyl (1R)			CH <sub>3</sub>	CH <sub>2</sub>	bond	-
45	H	n-Propyl	CN	H	CH <sub>3</sub>	CH <sub>2</sub>	bond	329 (M+H) <sup>+</sup>
46	H	n-Propyl (1R)	CN	H	CH <sub>3</sub>	CH <sub>2</sub>	bond	-







Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>7</sub>	R <sub>9</sub>	R <sub>10</sub>	Y	M	Mass Spec* (m/z)
47	H	<i>n</i> -Propyl (1S)	CN	H	CH <sub>3</sub>	CH <sub>2</sub>	bond	-
48	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	395 (M-H)-
49	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	420 (M+H) <sup>+</sup>
50	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	401 (M-H) <sup>-</sup>
51	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	435 (M+H) <sup>+</sup>
52	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	434 (M+H) <sup>+</sup>
53	H	<i>n</i> -Propyl (1R)	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	-
54	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	435 (M+H) <sup>+</sup>
55	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	434 (M+H) <sup>+</sup>
56	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	422 (M-H)-







Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>7</sub>	R <sub>9</sub>	R <sub>10</sub>	Y	M	Mass Spec* (m/z)
57	H	<i>n</i> -Propyl (1S)	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	424 (M+H) <sup>+</sup>
58	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	424 (M-H) <sup>-</sup>
59	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	435 (M+H) <sup>+</sup>
60	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	426 (M+H) <sup>+</sup>
61	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	464 (M+H) <sup>+</sup>
62	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	432 (M+H) <sup>+</sup>
63	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	432 (M+H) <sup>+</sup>
64	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	421 (M+H) <sup>+</sup>
65	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	463 (M+H) <sup>+</sup>


















Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>7</sub>	R <sub>9</sub>	R <sub>10</sub>	Y	M	Mass Spec* (m/z)
66	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	437 (M+H) <sup>+</sup>
67	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	448 (M+H) <sup>+</sup>
68	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	503 (M-H) <sup>-</sup>
69	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	433 (M-H) <sup>-</sup>
70	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	434 (M+H) <sup>+</sup>
71	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	436 (M+H) <sup>+</sup>
72	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	466 (M-H) <sup>-</sup>
73	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	488 (M+H) <sup>+</sup>










Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>7</sub>	R <sub>9</sub>	R <sub>10</sub>	Y	M	Mass Spec* (m/z)
74	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	424 (M-H) <sup>+</sup>
75	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	457 (M+H) <sup>+</sup>
76	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	409 (M+H) <sup>+</sup>
77	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	424 (M+H) <sup>+</sup>
78	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	437 (M+H) <sup>+</sup>
79	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	422 (M-H) <sup>+</sup>
80	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	436 (M-H) <sup>+</sup>










Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>7</sub>	R <sub>9</sub>	R <sub>10</sub>	Y	M	Mass Spec* (m/z)
81	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	488 (M-H) <sup>-</sup>
82	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	-
83	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	407 (M-H) <sup>-</sup>
84	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	451 (M+H) <sup>+</sup>
85	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	482 (M+H) <sup>+</sup>
86	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	464 (M+H) <sup>+</sup>

Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>7</sub>	R <sub>9</sub>	R <sub>10</sub>	Y	M	Mass Spec* (m/z)
87	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	483 (M+H) <sup>+</sup>
88	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	467 (M+H) <sup>+</sup>
89	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	450 (M-H) <sup>-</sup>
90	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	460 (M-H) <sup>-</sup>
91	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	476 (M-H) <sup>-</sup>
92	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	462 (M-H) <sup>-</sup>







Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>7</sub>	R <sub>9</sub>	R <sub>10</sub>	Y	M	Mass Spec* (m/z)
93	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	475 (M-H) <sup>+</sup>
94	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	462 (M-H) <sup>+</sup>
95	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	489 (M-H) <sup>+</sup>
96	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	511 (M-H) <sup>+</sup>
97	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	424 (M-H) <sup>+</sup>
98	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	383 (M-H) <sup>+</sup>
99	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	371 (M-H) <sup>+</sup>







Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>7</sub>	R <sub>9</sub>	R <sub>10</sub>	Y	M	Mass Spec* (m/z)
100	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	412 (M-H) <sup>+</sup>
101	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	424 (M-H) <sup>+</sup>
102	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	385 (M-H) <sup>+</sup>
103	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	398 (M-H) <sup>+</sup>
104	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	426 (M-H) <sup>+</sup>
105	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	426 (M-H) <sup>+</sup>
106	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	441 (M-H) <sup>+</sup>
107	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	473 (M-H) <sup>+</sup>







Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>7</sub>	R <sub>9</sub>	R <sub>10</sub>	Y	M	Mass Spec* (m/z)
108	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	425 (M-H) <sup>-</sup>
109	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	446 (M+H) <sup>+</sup>
110	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	421 (M-H) <sup>-</sup>
111	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	401 (M+H) <sup>+</sup>
112	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	415 (M+H) <sup>+</sup>
113	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	373 (M+H) <sup>+</sup>
114	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	476 (M+H) <sup>+</sup>
115	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	431 (M+H) <sup>+</sup>
116	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	423 (M-H) <sup>-</sup>

Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>7</sub>	R <sub>9</sub>	R <sub>10</sub>	Y	M	Mass Spec* (m/z)
117	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	463 (M+H) <sup>+</sup>
118	H	<i>n</i> -Propyl (1 <i>R</i> )	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	399 (M-H) <sup>-</sup>
119	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	399 (M-H) <sup>-</sup>
120	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	425 (M-H) <sup>-</sup>
121	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	399 (M-H) <sup>-</sup>
122	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	502 (M-H) <sup>-</sup>
123	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	474 (M-H) <sup>-</sup>
124	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	408 (M-H) <sup>-</sup>
125	H	<i>n</i> -Propyl (1 <i>R</i> )	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	422 (M-H) <sup>-</sup>



Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>7</sub>	R <sub>9</sub>	R <sub>10</sub>	Y	M	Mass Spec* (m/z)
126	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	489 (M-H) <sup>+</sup>
127	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	448 (M-H) <sup>+</sup>
128	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	421 (M-H) <sup>+</sup>
129	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	475 (M-H) <sup>+</sup>
130	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	461 (M-H) <sup>+</sup>
131	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	438 (M-H) <sup>+</sup>

Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>7</sub>	R <sub>9</sub>	R <sub>10</sub>	Y	M	Mass Spec* (m/z)
132	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	496 (M-H) <sup>+</sup>
133	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	441 (M+H) <sup>+</sup>
134	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	469 (M-H) <sup>+</sup>
135	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	452 (M+H) <sup>+</sup>
136	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	470 (M+H) <sup>+</sup>
137	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	437 (M+H) <sup>+</sup>

Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>7</sub>	R <sub>9</sub>	R <sub>10</sub>	Y	M	Mass Spec* (m/z)
138	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	463 (M-H) <sup>-</sup>
139	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	449 (M-H) <sup>-</sup>
140	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	449 (M-H) <sup>-</sup>
141	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	401 (M+H) <sup>+</sup>
142	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	415 (M-H) <sup>-</sup>
143	H	<i>n</i> -Propyl	CN		CH <sub>3</sub>	CH <sub>2</sub>	bond	447 (M-H) <sup>-</sup>

\* Mass Spectroscopy data is expressed as a mass to charge ratio (m/z) for either (M+1) or (M-1) molecular ion.

\*\* Compounds of Example No.s 45-47 are useful for making compounds of Formula I.

Example 1

(5-Cyano-7-hydroxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid

Example 2

(5-Cyano-7-methoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid

Example 3

(5-Cyano-7-ethoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid

Example 4

(5-Cyano-8-methyl-7-propoxymethyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid

Example 5

(5-Cyano-7-isopropoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid

Example 6

(5-Cyano-7-cyclobutoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid

Example 7

(5-Cyano-7-cyclohexyloxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid

Example 8

(5-Cyano-7-cyclopropylmethoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid

Example 9

(5-Cyano-7-cyclobutylmethoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid

Example 10

(5-Cyano-7-cyclopentylmethoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid

Example 11

(7-But-2-ynyloxymethyl-5-cyano-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid

Example 12

[5-Cyano-8-methyl-1-propyl-7-(tetrahydro-pyran-4-ylmethoxymethyl)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid

Example 13

(3'S, 1S\*) [5-Cyano-8-methyl-1-propyl-7-(tetrahydro-furan-3-yloxymethyl)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid

Example 14

(3'R, 1S\*) [5-Cyano-8-methyl-1-propyl-7-(tetrahydro-furan-3-yloxymethyl)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid

Example 15

(7-Benzylloxymethyl-5-cyano-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid

Example 16

[7-(Benzo[1,3]dioxol-5-ylmethoxymethyl)-5-cyano-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid

Example 17

[5-Cyano-7-(2,4-dimethyl-benzyloxymethyl)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid

Example 18

[5-Cyano-8-methyl-1-propyl-7-(thiophen-3-ylmethoxymethyl)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid

Example 19

[5-Cyano-7-(2,4-dimethyl-thiazol-5-ylmethoxymethyl)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid

Example 20

(5-Cyano-8-methyl-7-phenoxyethyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid

Example 21

[5-Cyano-7-(3-fluoro-phenoxyethyl)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid

Example 22

(5-Cyano-7-cyclopropylmethoxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid

Example 23

(R)-[5-Cyano-8-methyl-7-(5-methyl-isoxazol-3-ylmethoxy)-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid

Example 24

[5-Cyano-8-methyl-1-propyl-7-(pyridin-4-ylmethoxy)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid

Example 25

[5-Cyano-7-(1,5-dimethyl-1H-pyrazol-3-ylmethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid

Example 26

(R)-[5-Cyano-7-(2-isopropoxy-ethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid

Example 27

(R)-[5-Cyano-7-(3-methoxy-propoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid

Example 28

(1R, 2'R)-[5-Cyano-7-(2-methoxy-propoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid

Example 29

[5-Cyano-8-methyl-7-(5-methyl-[1,3,4]thiadiazol-2-ylmethoxy)-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid

Example 30

(R)-[5-Cyano-7-(5-dimethylamino-[1,2,4]thiadiazol-3-ylmethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid

Example 31

5-Cyano-7-(2-methoxy-ethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indole-1-carboxylic acid

Example 32

5-Cyano-8-methyl-7-(5-methyl-isoxazol-3-ylmethoxy)-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indole-1-carboxylic acid

Example 33

(1*R*\*,10*S*)- [1-sec-Butyl-5-cyano-7-(2-ethoxy-ethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid

Example 34

(1*R*\*,10*S*)- [1-sec-Butyl-5-cyano-7-(2-isopropoxy-ethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid

Example 35

(1*R*\*,10*S*)- [1-sec-Butyl-5-cyano-7-(5-dimethylamino-[1,2,4]thiadiazol-3-ylmethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid

Example 36


(1*R*\*,10*S*)-[1-sec-Butyl-5-cyano-7-(1,5-dimethyl-1*H*-pyrazol-3-ylmethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-*b*]indol-1-yl]-acetic acid



CLAIMS

What is claimed:

1. A compound having the formula:



wherein:

R<sub>1</sub> is hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, or an arylalkyl or an alkylaryl of 7 to 12 carbon atoms;

R<sub>2</sub> is hydrogen, a straight chain alkyl of 1 to 12 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, an alkoxyalkyl of 2 to 12 carbon atoms, an arylalkyl or alkylaryl of 7 to 12 carbon atoms, a cyanoalkyl of 1 to 8 carbon atoms, an alkylthioalkyl of 2 to 16 carbon atoms, a cycloalkyl-alkyl of 4 to 24 carbon atoms, a substituted or unsubstituted aryl, or a heteroaryl;

R<sub>3</sub> - R<sub>6</sub> are independently hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, a substituted or unsubstituted aryl, furanylmethyl, arylalkyl or

alkylaryl of 7 to 12 carbon atoms, alkynyl of 2 to 7 carbon atoms, or R<sub>5</sub> and R<sub>6</sub> together with the ring carbon atom to which they are attached form a carbonyl group;

R<sub>7</sub>- R<sub>8</sub> and R<sub>10</sub> are independently hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, furanylmethyl, arylalkyl or alkylaryl of 7 to 12 carbon atoms, alkynyl of 2 to 7 carbon atoms, phenylalkynyl, alkoxy of 1 to 8 carbon atoms, arylalkoxy of 7 to 12 carbon atoms, alkylthio of 1 to 8 carbon atoms, trifluoromethoxy, trifluoroethoxy, trifluoromethylthio, trifluoroethylthio, acyl of 1 to 6 carbon atoms, COOH, COO-alkyl, CONR<sub>11</sub>R<sub>12</sub>, F, Cl, Br, I, CN, CF<sub>3</sub>, NO<sub>2</sub>, alkylsulfinyl of 1 to 8 carbon atoms, alkylsulfonyl of 1 to 6 carbon atoms, pyrrolidinyl, or thiazolidinyl;

R<sub>9</sub> is hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, a cycloalkyl of 3 to 12 carbon atoms, a cycloalkyl-alkyl of 4 to 24 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, an alkoxyalkyl of 2 to 12 carbon atoms, an alkoxyalkoxyalkyl of 3 to 18 carbon atoms, an arylalkoxyalkyl of 3 to 18 carbon atoms, a cycloalkylalkoxyalkyl of 3 to 18 carbon atoms, an aryloxyalkyl of 3 to 18 carbon atoms, a heteroaryloxyalkyl of 3 to 18 carbon atoms, an arylthioalkyl of 3 to 18 carbon atoms, a heteroarylthioalkyl of 3 to 18 carbon atoms, a hydroxyalkyl of 1 to 12 carbon atoms, an alkoxyiminoalkyl of 2 to 16 carbon atoms, an alkylthioalkyl of 2 to 16 carbon atoms, an alkylsulfonylalkyl group of 2 to 16 carbon atoms, a monoalkylaminoalkyl of 2 to 16 carbon atoms, a dialkylaminoalkyl of 3 to 16 carbon atoms, a substituted dialkylaminoalkyl of 3 to 16 carbon atoms, a substituted or unsubstituted aryl, arylalkyl of 7 to 12 carbon atoms, a substituted or unsubstituted heteroaryl of 7 to 12 carbon atoms, a substituted or unsubstituted heteroarylalkyl, a substituted or unsubstituted heterocyclic group, and a heterocycle-alkyl;

R<sub>11</sub> - R<sub>12</sub> are independently H, straight chain alkyl of 1 to 8 carbon atoms, branched alkyl of 3 to 12 carbon atoms, cycloalkyl of 3 to 12 carbon atoms, a substituted or unsubstituted aryl or heteroaryl;

M is a bond, CH<sub>2</sub>, or CH<sub>2</sub>CH<sub>2</sub>, with the proviso that when M is a bond, then R<sub>9</sub> is other than a hydroxyl, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, or an arylalkyl;

Y is a bond, CH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>, aryl, or R<sub>2</sub> and Y together with the ring carbon atom to which they are attached may additionally form a spirocyclic cycloalkyl ring of 3 to 8 carbon atoms; or

a crystalline form or a pharmaceutically acceptable salt thereof.

2. The compound according to claim 1 wherein

R<sub>1</sub> is hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, or an arylalkyl or an alkylaryl of 7 to 12 carbon atoms;

R<sub>2</sub> is hydrogen, a straight chain alkyl of 1 to 12 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, an alkoxyalkyl of 2 to 12 carbon atoms, an arylalkyl or alkylaryl of 7 to 12 carbon atoms, a cyanoalkyl of 1 to 8 carbon atoms, an alkylthioalkyl of 2 to 16 carbon atoms, a cycloalkyl-alkyl of 4 to 24 carbon atoms, a substituted or unsubstituted aryl, or a heteroaryl;

R<sub>3</sub> - R<sub>6</sub> are independently hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, a substituted or unsubstituted aryl, furanylmethyl, arylalkyl or alkylaryl of 7 to 12 carbon atoms, alkynyl of 2 to 7 carbon atoms, or R<sub>5</sub> and R<sub>6</sub> together with the ring carbon atom to which they are attached form a carbonyl group;

R<sub>7</sub>- R<sub>8</sub> and R<sub>10</sub> are independently hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, furanylmethyl, arylalkyl or alkylaryl of 7 to 12 carbon atoms, alkynyl of 2 to 7 carbon atoms, phenylalkynyl, alkoxy of 1 to 8 carbon atoms, arylalkoxy of 7 to 12 carbon atoms, alkylthio of 1 to 8 carbon atoms, trifluoromethoxy, trifluoroethoxy, trifluoromethylthio, trifluoroethylthio, acyl of 1 to 6 carbon atoms, COOH, COO-alkyl, CONR<sub>11</sub>R<sub>12</sub>, F, Cl, Br, I, CN, CF<sub>3</sub>, NO<sub>2</sub>, alkylsulfinyl of 1 to 8 carbon atoms, alkylsulfonyl of 1 to 6 carbon atoms, pyrrolidinyl, or thiazolidinyl;

R<sub>9</sub> is hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, a cycloalkyl of 3 to 12 carbon atoms, a cycloalkyl-alkyl of 4 to 24 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, an alkoxyalkyl of 2 to 12 carbon atoms, an alkoxyalkoxyalkyl of 3 to 18 carbon atoms, an arylalkoxyalkyl of 3 to 18 carbon atoms, a cycloalkylalkoxyalkyl of 3 to 18 carbon atoms, an aryloxyalkyl of 3 to 18 carbon atoms, a heteroaryloxyalkyl of 3 to 18 carbon atoms, an arylthioalkyl of 3 to 18 carbon atoms, a heteroarylthioalkyl of 3 to 18 carbon atoms, a hydroxyalkyl of 1 to 12 carbon atoms, an alkoxyiminoalkyl of 2 to 16 carbon atoms, an alkylthioalkyl of 2 to 16 carbon atoms, an alkylsulfonylalkyl group of 2 to 16 carbon atoms, a monoalkylaminoalkyl of 2 to 16 carbon atoms, a dialkylaminoalkyl of 3 to 16 carbon atoms, a substituted dialkylaminoalkyl of 3 to 16 carbon atoms, a substituted or unsubstituted aryl, arylalkyl of 7 to 12 carbon atoms, a substituted or unsubstituted heteroaryl of 7 to 12 carbon atoms, a substituted or unsubstituted heteroarylalkyl, a substituted or unsubstituted heterocyclic group, and a heterocycle-alkyl;

R<sub>11</sub> - R<sub>12</sub> are independently H, straight chain alkyl of 1 to 8 carbon atoms, branched alkyl of 3 to 12 carbon atoms, cycloalkyl of 3 to 12 carbon atoms, a substituted or unsubstituted aryl or heteroaryl;

M is CH<sub>2</sub> or CH<sub>2</sub>CH<sub>2</sub>;

Y is a bond, CH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>, aryl, or R<sub>2</sub> and Y together with the ring carbon atom to which they are attached may additionally form a spirocyclic cycloalkyl ring of 3 to 8 carbon atoms; or

a crystalline form or a pharmaceutically acceptable salt thereof.

3. The compound according to claim 1 wherein:

R<sub>1</sub> is hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, or an arylalkyl or an alkylaryl of 7 to 12 carbon atoms;

R<sub>2</sub> is hydrogen, a straight chain alkyl of 1 to 12 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms; an alkynyl of 2 to 7 carbon atoms, an alkoxyalkyl of 2 to 12 carbon atoms, an arylalkyl or alkylaryl of 7 to 12 carbon atoms, a cyanoalkyl of 1 to 8 carbon atoms, an alkylthioalkyl of 2 to 16 carbon atoms, a cycloalkyl-alkyl of 4 to 24 carbon atoms, a substituted or unsubstituted aryl, or a heteroaryl;

R<sub>3</sub> - R<sub>6</sub> are independently hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, a substituted or unsubstituted aryl, furanylmethyl, arylalkyl or alkylaryl of 7 to 12 carbon atoms, alkynyl of 2 to 7 carbon atoms, or R<sub>5</sub> and R<sub>6</sub> together with the ring carbon atom to which they are attached form a carbonyl group;

R<sub>7</sub>- R<sub>8</sub> and R<sub>10</sub> are independently hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, furanylmethyl, arylalkyl or alkylaryl of 7 to 12 carbon atoms, alkynyl of 2 to 7 carbon atoms, phenylalkynyl, alkoxy of 1 to 8 carbon atoms, arylalkoxy

of 7 to 12 carbon atoms, alkylthio of 1 to 8 carbon atoms, trifluoromethoxy, trifluoroethoxy, trifluoromethylthio, trifluoroethylthio, acyl of 1 to 6 carbon atoms, COOH, COO-alkyl, CONR<sub>11</sub>R<sub>12</sub>, F, Cl, Br, I, CN, CF<sub>3</sub>, NO<sub>2</sub>, alkylsulfinyl of 1 to 8 carbon atoms, alkylsulfonyl of 1 to 6 carbon atoms, pyrrolidinyl, or thiazolidinyl;

R<sub>9</sub> is a cycloalkyl of 3 to 12 carbon atoms, a cycloalkyl-alkyl of 4 to 24 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, an alkoxyalkyl of 2 to 12 carbon atoms, an alkoxyalkoxyalkyl of 3 to 18 carbon atoms, an arylalkoxyalkyl of 3 to 18 carbon atoms, a cycloalkylalkoxyalkyl of 3 to 18 carbon atoms, an aryloxyalkyl of 3 to 18 carbon atoms, a heteroaryloxyalkyl of 3 to 18 carbon atoms, an arylthioalkyl of 3 to 18 carbon atoms, a heteroarylthioalkyl of 3 to 18 carbon atoms, a hydroxyalkyl of 1 to 12 carbon atoms, an alkoxyiminoalkyl of 2 to 16 carbon atoms, an alkylthioalkyl of 2 to 16 carbon atoms, an alkylsulfonylalkyl group of 2 to 16 carbon atoms, a monoalkylaminoalkyl of 2 to 16 carbon atoms, a dialkylaminoalkyl of 3 to 16 carbon atoms, a substituted dialkylaminoalkyl of 3 to 16 carbon atoms, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl of 7 to 12 carbon atoms, a substituted or unsubstituted heteroarylalkyl, a substituted or unsubstituted heterocyclic group, and a heterocycle-alkyl;

R<sub>11</sub> - R<sub>12</sub> are independently H, straight chain alkyl of 1 to 8 carbon atoms, branched alkyl of 3 to 12 carbon atoms, cycloalkyl of 3 to 12 carbon atoms, a substituted or unsubstituted aryl or heteroaryl;

M is a bond;

Y is a bond, CH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>, aryl, or R<sub>2</sub> and Y together with the ring carbon atom to which they are attached may additionally form a spirocyclic cycloalkyl ring of 3 to 8 carbon atoms; or

a crystalline form or a pharmaceutically acceptable salt thereof.

## 4. The compound according to claim 1 selected from the group consisting of:

- (5-cyano-7-hydroxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
- (5-cyano-7-methoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
- (5-cyano-7-ethoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
- (5-cyano-8-methyl-7-propoxymethyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
- (5-cyano-7-isopropoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
- (5-cyano-7-cyclobutoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
- (5-cyano-7-cyclohexyloxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
- (5-cyano-7-cyclopropylmethoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
- (5-cyano-7-cyclobutylmethoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
- (5-cyano-7-cyclopentylmethoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
- (7-but-2-ynyloxymethyl-5-cyano-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
- [5-cyano-8-methyl-1-propyl-7-(tetrahydro-pyran-4-ylmethoxymethyl)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;
- (3'S, 1S\*) [5-cyano-8-methyl-1-propyl-7-(tetrahydro-furan-3-yloxymethyl)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;
- (3'R, 1S\*) [5-cyano-8-methyl-1-propyl-7-(tetrahydro-furan-3-yloxymethyl)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(7-benzyloxymethyl-5-cyano-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;

[7-(benzo[1,3]dioxol-5-ylmethoxymethyl)-5-cyano-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

[5-cyano-7-(2,4-dimethyl-benzyloxymethyl)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

[5-cyano-8-methyl-1-propyl-7-(thiophen-3-ylmethoxymethyl)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

[5-cyano-7-(2,4-dimethyl-thiazol-5-ylmethoxymethyl)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(5-cyano-8-methyl-7-phenoxy-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;

[5-cyano-7-(3-fluoro-phenoxy-methyl)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(5-cyano-7-cyclopropylmethoxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;

(R)-[5-cyano-8-methyl-7-(5-methyl-isoxazol-3-ylmethoxy)-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

[5-cyano-8-methyl-1-propyl-7-(pyridin-4-ylmethoxy)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

[5-cyano-7-(1,5-dimethyl-1H-pyrazol-3-ylmethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(R)-[5-cyano-7-(2-isopropoxy-ethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(R)-[5-cyano-7-(3-methoxy-propoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(1R, 2'R)-[5-cyano-7-(2-methoxy-propoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

[5-cyano-8-methyl-7-(5-methyl-[1,3,4]thiadiazol-2-ylmethoxy)-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;



(R)-[5-cyano-7-(5-dimethylamino-[1,2,4]thiadiazol-3-ylmethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

5-cyano-7-(2-methoxy-ethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4,-b]indole-1-carboxylic acid;

5-cyano-8-methyl-7-(5-methyl-isoxazol-3-ylmethoxy)-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4,-b]indole-1-carboxylic acid;


(1*R*\*,10*S*)-[1-sec-butyl-5-cyano-7-(2-ethoxy-ethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(1*R*\*,10*S*)- [1-sec-butyl-5-cyano-7-(2-isopropoxy-ethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;


(1*R*\*,10*S*)- [1-sec-butyl-5-cyano-7-(5-dimethylamino-[1,2,4]thiadiazol-3-ylmethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid; and

(1*R*\*,10*S*)-[1-sec-butyl-5-cyano-7-(1,5-dimethyl-1*H*-pyrazol-3-ylmethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid.

5. The compound according to claim 1 having the formula:



6. A pharmaceutical composition comprising a compound of a formula:



wherein:

R<sub>1</sub> is hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, or an arylalkyl or an alkylaryl of 7 to 12 carbon atoms;

R<sub>2</sub> is hydrogen, a straight chain alkyl of 1 to 12 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, an alkoxyalkyl of 2 to 12 carbon atoms, an arylalkyl or alkylaryl of 7 to 12 carbon atoms, a cyanoalkyl of 1 to 8 carbon atoms, an alkylthioalkyl of 2 to 16 carbon atoms, a cycloalkyl-alkyl of 4 to 24 carbon atoms, a substituted or unsubstituted aryl, or a heteroaryl;

R<sub>3</sub> - R<sub>6</sub> are independently hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, a substituted or unsubstituted aryl, furanylmethyl, arylalkyl or alkylaryl of 7 to 12 carbon atoms, alkynyl of 2 to 7 carbon atoms, or R<sub>5</sub> and R<sub>6</sub> together with the ring carbon atom to which they are attached form a carbonyl group;

R<sub>7</sub>- R<sub>8</sub> and R<sub>10</sub> are independently hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, furanylmethyl, arylalkyl or alkylaryl of 7 to 12 carbon atoms, alkynyl of 2 to 7 carbon atoms, phenylalkynyl, alkoxy of 1 to 8 carbon atoms, arylalkoxy

of 7 to 12 carbon atoms, alkylthio of 1 to 8 carbon atoms, trifluoromethoxy, trifluoroethoxy, trifluoromethylthio, trifluoroethylthio, acyl of 1 to 6 carbon atoms, COOH, COO-alkyl, CONR<sub>11</sub>R<sub>12</sub>, F, Cl, Br, I, CN, CF<sub>3</sub>, NO<sub>2</sub>, alkylsulfinyl of 1 to 8 carbon atoms, alkylsulfonyl of 1 to 6 carbon atoms, pyrrolidinyl, or thiazolidinyl;

R<sub>9</sub> is hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, a cycloalkyl of 3 to 12 carbon atoms, a cycloalkyl-alkyl of 4 to 24 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, an alkoxyalkyl of 2 to 12 carbon atoms, an alkoxyalkoxyalkyl of 3 to 18 carbon atoms, an arylalkoxyalkyl of 3 to 18 carbon atoms, a cycloalkylalkoxyalkyl of 3 to 18 carbon atoms, an aryloxyalkyl of 3 to 18 carbon atoms, a heteroaryloxyalkyl of 3 to 18 carbon atoms, an arylthioalkyl of 3 to 18 carbon atoms, a heteroarylthioalkyl of 3 to 18 carbon atoms, a hydroxyalkyl of 1 to 12 carbon atoms, an alkoxyiminoalkyl of 2 to 16 carbon atoms, an alkylthioalkyl of 2 to 16 carbon atoms, an alkylsulfonylalkyl group of 2 to 16 carbon atoms, a monoalkylaminoalkyl of 2 to 16 carbon atoms, a dialkylaminoalkyl of 3 to 16 carbon atoms, a substituted dialkylaminoalkyl of 3 to 16 carbon atoms, a substituted or unsubstituted aryl, arylalkyl of 7 to 12 carbon atoms, a substituted or unsubstituted heteroaryl of 7 to 12 carbon atoms, a substituted or unsubstituted heteroarylalkyl, a substituted or unsubstituted heterocyclic group, and a heterocycle-alkyl;


R<sub>11</sub> - R<sub>12</sub> are independently H, straight chain alkyl of 1 to 8 carbon atoms, branched alkyl of 3 to 12 carbon atoms, cycloalkyl of 3 to 12 carbon atoms, a substituted or unsubstituted aryl or heteroaryl;

M is a bond, CH<sub>2</sub>, or CH<sub>2</sub>CH<sub>2</sub>, with the proviso that when M is a bond, then R<sub>9</sub> is other than a hydroxyl, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, or an arylalkyl;

Y is a bond, CH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>, aryl, or R<sub>2</sub> and Y together with the ring carbon atom to which they are attached may additionally form a spirocyclic cycloalkyl ring of 3 to 8 carbon atoms;

or a crystalline form or a pharmaceutically acceptable salt thereof; and a pharmaceutically acceptable carrier.

7. A method of treating or preventing a Hepatitis C viral infection in a mammal comprising providing the mammal with an effective amount of a compound of a formula:



wherein:

R<sub>1</sub> is hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, or an arylalkyl or an alkylaryl of 7 to 12 carbon atoms;

R<sub>2</sub> is hydrogen, a straight chain alkyl of 1 to 12 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, an alkoxyalkyl of 2 to 12 carbon atoms, an arylalkyl or alkylaryl of 7 to 12 carbon atoms, a cyanoalkyl of 1 to 8 carbon atoms, an alkylthioalkyl of 2 to 16 carbon atoms, a cycloalkyl-alkyl of 4 to 24 carbon atoms, a substituted or unsubstituted aryl, or a heteroaryl;

R<sub>3</sub> - R<sub>6</sub> are independently hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, a substituted or unsubstituted aryl, furanylmethyl, arylalkyl or

alkylaryl of 7 to 12 carbon atoms, alkynyl of 2 to 7 carbon atoms, or R<sub>5</sub> and R<sub>6</sub> together with the ring carbon atom to which they are attached form a carbonyl group;

R<sub>7</sub>- R<sub>8</sub> and R<sub>10</sub> are independently hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, furanylmethyl, arylalkyl or alkylaryl of 7 to 12 carbon atoms, alkynyl of 2 to 7 carbon atoms, phenylalkynyl, alkoxy of 1 to 8 carbon atoms, arylalkoxy of 7 to 12 carbon atoms, alkylthio of 1 to 8 carbon atoms, trifluoromethoxy, trifluoroethoxy, trifluoromethylthio, trifluoroethylthio, acyl of 1 to 6 carbon atoms, COOH, COO-alkyl, CONR<sub>11</sub>R<sub>12</sub>, F, Cl, Br, I, CN, CF<sub>3</sub>, NO<sub>2</sub>, alkylsulfinyl of 1 to 8 carbon atoms, alkylsulfonyl of 1 to 6 carbon atoms, pyrrolidinyl, or thiazolidinyl;

R<sub>9</sub> is hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, a cycloalkyl of 3 to 12 carbon atoms, a cycloalkyl-alkyl of 4 to 24 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, an alkoxyalkyl of 2 to 12 carbon atoms, an alkoxyalkoxyalkyl of 3 to 18 carbon atoms, an arylalkoxyalkyl of 3 to 18 carbon atoms, a cycloalkylalkoxyalkyl of 3 to 18 carbon atoms, an aryloxyalkyl of 3 to 18 carbon atoms, a heteroaryloxyalkyl of 3 to 18 carbon atoms, an arylthioalkyl of 3 to 18 carbon atoms, a heteroarylthioalkyl of 3 to 18 carbon atoms, a hydroxyalkyl of 1 to 12 carbon atoms, an alkoxyiminoalkyl of 2 to 16 carbon atoms, an alkylthioalkyl of 2 to 16 carbon atoms, an alkylsulfonylalkyl group of 2 to 16 carbon atoms, a monoalkylaminoalkyl of 2 to 16 carbon atoms, a dialkylaminoalkyl of 3 to 16 carbon atoms, a substituted dialkylaminoalkyl of 3 to 16 carbon atoms, a substituted or unsubstituted aryl, arylalkyl of 7 to 12 carbon atoms, a substituted or unsubstituted heteroaryl of 7 to 12 carbon atoms, a substituted or unsubstituted heteroarylalkyl, a substituted or unsubstituted heterocyclic group, and a heterocycle-alkyl;

R<sub>11</sub> - R<sub>12</sub> are independently H, straight chain alkyl of 1 to 8 carbon atoms, branched alkyl of 3 to 12 carbon atoms, cycloalkyl of 3 to 12 carbon atoms, a substituted or unsubstituted aryl or heteroaryl;

M is a bond, CH<sub>2</sub>, or CH<sub>2</sub>CH<sub>2</sub>, with the proviso that when M is a bond, then R<sub>9</sub> is other than a hydroxyl, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, or an arylalkyl;

Y is a bond, CH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>, aryl, or R<sub>2</sub> and Y together with the ring carbon atom to which they are attached may additionally form a spirocyclic cycloalkyl ring of 3 to 8 carbon atoms; or

a crystalline form or a pharmaceutically acceptable salt thereof.

8. The method of Claim 7 wherein the compound is selected from the group consisting of
  - (5-cyano-7-hydroxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
  - (5-cyano-7-methoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
  - (5-cyano-7-ethoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
  - (5-cyano-8-methyl-7-propoxymethyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
  - (5-cyano-7-isopropoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
  - (5-cyano-7-cyclobutoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
  - (5-cyano-7-cyclohexyloxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
  - (5-cyano-7-cyclopropylmethoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;

(5-cyano-7-cyclobutylmethoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;

(5-cyano-7-cyclopentylmethoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;

(7-but-2-ynyloxymethyl-5-cyano-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;

[5-cyano-8-methyl-1-propyl-7-(tetrahydro-pyran-4-ylmethoxymethyl)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(3'S, 1S\*) [5-cyano-8-methyl-1-propyl-7-(tetrahydro-furan-3-yloxymethyl)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(3'R, 1S\*) [5-cyano-8-methyl-1-propyl-7-(tetrahydro-furan-3-yloxymethyl)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(7-benzyloxymethyl-5-cyano-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;

[7-(benzo[1,3]dioxol-5-ylmethoxymethyl)-5-cyano-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

[5-cyano-7-(2,4-dimethyl-benzyloxymethyl)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

[5-cyano-8-methyl-1-propyl-7-(thiophen-3-ylmethoxymethyl)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

[5-cyano-7-(2,4-dimethyl-thiazol-5-ylmethoxymethyl)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(5-cyano-8-methyl-7-phenoxy-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;

[5-cyano-7-(3-fluoro-phenoxy-methyl)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(5-cyano-7-cyclopropylmethoxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;

(R)-[5-cyano-8-methyl-7-(5-methyl-isoxazol-3-ylmethoxy)-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

[5-cyano-8-methyl-1-propyl-7-(pyridin-4-ylmethoxy)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

[5-cyano-7-(1,5-dimethyl-1H-pyrazol-3-ylmethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(R)-[5-cyano-7-(2-isopropoxy-ethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(R)-[5-cyano-7-(3-methoxy-propoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(1R, 2'R)-[5-cyano-7-(2-methoxy-propoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

[5-cyano-8-methyl-7-(5-methyl-[1,3,4]thiadiazol-2-ylmethoxy)-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(R)-[5-cyano-7-(5-dimethylamino-[1,2,4]thiadiazol-3-ylmethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

5-cyano-7-(2-methoxy-ethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indole-1-carboxylic acid;

5-cyano-8-methyl-7-(5-methyl-isoxazol-3-ylmethoxy)-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indole-1-carboxylic acid;

(1R\*,10S)-[1-sec-butyl-5-cyano-7-(2-ethoxy-ethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;


(1R\*,10S)- [1-sec-butyl-5-cyano-7-(2-isopropoxy-ethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(1R\*,10S)- [1-sec-butyl-5-cyano-7-(5-dimethylamino-[1,2,4]thiadiazol-3-ylmethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid; and


(1R\*,10S)-[1-sec-butyl-5-cyano-7-(1,5-dimethyl-1H-pyrazol-3-ylmethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid.

9. The method of Claim 7 wherein the compound of the formula has a ratio of Isomer A to Isomer B of greater than 1:1, wherein Isomer A and Isomer B have the respective formulas:






Isomer A



Isomer B.

10. The method of Claim 7 herein the compound of the formula is 100% Isomer A.
11. The method of Claim 7 wherein the compound of the formula has a ratio of Isomer A to Isomer B of at least 9:1.
12. The method of Claim 7 wherein the compound of the formula has a ratio of Isomer A to Isomer B of at least 8:1.
13. The method of Claim 7 wherein the compound of the formula has a ratio of Isomer A to Isomer B of at least 7:1.

14. A method of inhibiting replication of a Hepatitis C virus comprising contacting the Hepatitis C virus with a compound of a formula:



wherein:

R<sub>1</sub> is hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, or an arylalkyl or an alkylaryl of 7 to 12 carbon atoms;

R<sub>2</sub> is hydrogen, a straight chain alkyl of 1 to 12 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, an alkoxyalkyl of 2 to 12 carbon atoms, an arylalkyl or alkylaryl of 7 to 12 carbon atoms, a cyanoalkyl of 1 to 8 carbon atoms, an alkylthioalkyl of 2 to 16 carbon atoms, a cycloalkyl-alkyl of 4 to 24 carbon atoms, a substituted or unsubstituted aryl, or a heteroaryl;

R<sub>3</sub> - R<sub>6</sub> are independently hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbon atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, a substituted or unsubstituted aryl, furanylmethyl, arylalkyl or alkylaryl of 7 to 12 carbon atoms, alkynyl of 2 to 7 carbon atoms, or R<sub>5</sub> and R<sub>6</sub> together with the ring carbon atom to which they are attached form a carbonyl group;

R<sub>7</sub>- R<sub>8</sub> and R<sub>10</sub> are independently hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, a cycloalkyl of 3 to 12 carbon atoms, an alkenyl of 2 to 7 carbon atoms, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, furanylmethyl, arylalkyl or alkylaryl of 7 to 12 carbon atoms, alkynyl of 2 to 7 carbon atoms, phenylalkynyl, alkoxy of 1 to 8 carbon atoms, arylalkoxy of 7 to 12 carbon atoms, alkylthio of 1 to 8 carbon atoms, trifluoromethoxy, trifluoroethoxy, trifluoromethylthio, trifluoroethylthio, acyl of 1 to 6 carbon atoms, COOH, COO-alkyl, CONR<sub>11</sub>R<sub>12</sub>, F, Cl, Br, I, CN, CF<sub>3</sub>, NO<sub>2</sub>, alkylsulfinyl of 1 to 8 carbon atoms, alkylsulfonyl of 1 to 6 carbon atoms, pyrrolidinyl, or thiazolidinyl;

R<sub>9</sub> is hydrogen, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, a cycloalkyl of 3 to 12 carbon atoms, a cycloalkyl-alkyl of 4 to 24 carbon atoms, an alkenyl of 2 to 7 carbon atoms, an alkynyl of 2 to 7 carbon atoms, an alkoxyalkyl of 2 to 12 carbon atoms, an alkoxyalkoxyalkyl of 3 to 18 carbon atoms, an arylalkoxyalkyl of 3 to 18 carbon atoms, a cycloalkylalkoxyalkyl of 3 to 18 carbon atoms, an aryloxyalkyl of 3 to 18 carbon atoms, a heteroaryloxyalkyl of 3 to 18 carbon atoms, an arylthioalkyl of 3 to 18 carbon atoms, a heteroarylthioalkyl of 3 to 18 carbon atoms, a hydroxyalkyl of 1 to 12 carbon atoms, an alkoxyiminoalkyl of 2 to 16 carbon atoms, an alkylthioalkyl of 2 to 16 carbon atoms, an alkylsulfonylalkyl group of 2 to 16 carbon atoms, a monoalkylaminoalkyl of 2 to 16 carbon atoms, a dialkylaminoalkyl of 3 to 16 carbon atoms, a substituted dialkylaminoalkyl of 3 to 16 carbon atoms, a substituted or unsubstituted aryl, arylalkyl of 7 to 12 carbon atoms, a substituted or unsubstituted heteroaryl of 7 to 12 carbon atoms, a substituted or unsubstituted heteroarylalkyl, a substituted or unsubstituted heterocyclic group, and a heterocycle-alkyl;


R<sub>11</sub> - R<sub>12</sub> are independently H, straight chain alkyl of 1 to 8 carbon atoms, branched alkyl of 3 to 12 carbon atoms, cycloalkyl of 3 to 12 carbon atoms, a substituted or unsubstituted aryl or heteroaryl;

M is a bond, CH<sub>2</sub>, or CH<sub>2</sub>CH<sub>2</sub>, with the proviso that when M is a bond, then R<sub>9</sub> is other than a hydroxyl, a straight chain alkyl of 1 to 8 carbon atoms, a branched alkyl of 3 to 12 carbons atoms, or an arylalkyl;


Y is a bond, CH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>, aryl, or R<sub>2</sub> and Y together with the ring carbon atom to which they are attached may additionally form a spirocyclic cycloalkyl ring of 3 to 8 carbon atoms; or

a crystalline form or a pharmaceutically acceptable salt thereof.

15. The method of Claim 14 wherein the compound of the formula has a ratio of Isomer A to Isomer B of greater than 1:1, wherein Isomer A and Isomer B have the respective formulas:



Isomer A



Isomer B.

16. The method of Claim 14 wherein the compound of the formula has a ratio of R-enantiomer to S-enantiomer of greater than 1:1.
17. The method of Claim 14 wherein the compound of the formula is 100% Isomer A.
18. The method of Claim 14 wherein the compound of the formula has a ratio of Isomer A to Isomer B of at least 9:1.
19. The method of Claim 14 wherein the compound of the formula has a ratio of Isomer A to Isomer B of at least 8:1.
20. The method of Claim 14 wherein the compound of the formula has a ratio of Isomer A to Isomer B of at least 7:1.
21. The method of Claim 14 wherein the compound is selected from:
  - (5-cyano-7-hydroxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
  - (5-cyano-7-methoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
  - (5-cyano-7-ethoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
  - (5-cyano-8-methyl-7-propoxymethyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
  - (5-cyano-7-isopropoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;
  - (5-cyano-7-cyclobutoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;

(5-cyano-7-cyclohexyloxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;

(5-cyano-7-cyclopropylmethoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;

(5-cyano-7-cyclobutylmethoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;

(5-cyano-7-cyclopentylmethoxymethyl-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;

(7-but-2-ynyloxymethyl-5-cyano-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;

[5-cyano-8-methyl-1-propyl-7-(tetrahydro-pyran-4-ylmethoxymethyl)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(3'S, 1S\*) [5-cyano-8-methyl-1-propyl-7-(tetrahydro-furan-3-yloxymethyl)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(3'R, 1S\*) [5-cyano-8-methyl-1-propyl-7-(tetrahydro-furan-3-yloxymethyl)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(7-benzyloxymethyl-5-cyano-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;

[7-(benzo[1,3]dioxol-5-ylmethoxymethyl)-5-cyano-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

[5-cyano-7-(2,4-dimethyl-benzyloxymethyl)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

[5-cyano-8-methyl-1-propyl-7-(thiophen-3-ylmethoxymethyl)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

[5-cyano-7-(2,4-dimethyl-thiazol-5-ylmethoxymethyl)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(5-cyano-8-methyl-7-phenoxyethyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;

[5-cyano-7-(3-fluoro-phenoxyethyl)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(5-cyano-7-cyclopropylmethoxy-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl)-acetic acid;

(R)-[5-cyano-8-methyl-7-(5-methyl-isoxazol-3-ylmethoxy)-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

[5-cyano-8-methyl-1-propyl-7-(pyridin-4-ylmethoxy)-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

[5-cyano-7-(1,5-dimethyl-1H-pyrazol-3-ylmethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(R)-[5-cyano-7-(2-isopropoxy-ethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(R)-[5-cyano-7-(3-methoxy-propoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(1R, 2'R)-[5-cyano-7-(2-methoxy-propoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

[5-cyano-8-methyl-7-(5-methyl-[1,3,4]thiadiazol-2-ylmethoxy)-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(R)-[5-cyano-7-(5-dimethylamino-[1,2,4]thiadiazol-3-ylmethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

5-cyano-7-(2-methoxy-ethoxy)-8-methyl-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indole-1-carboxylic acid;

5-cyano-8-methyl-7-(5-methyl-isoxazol-3-ylmethoxy)-1-propyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indole-1-carboxylic acid;

(1R\*,10S)-[1-sec-butyl-5-cyano-7-(2-ethoxy-ethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(1R\*,10S)-[1-sec-butyl-5-cyano-7-(2-isopropoxy-ethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid;

(1R\*,10S)-[1-sec-butyl-5-cyano-7-(5-dimethylamino-[1,2,4]thiadiazol-3-ylmethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid; and

(1R\*,10S)-[1-sec-butyl-5-cyano-7-(1,5-dimethyl-1H-pyrazol-3-ylmethoxy)-8-methyl-1,3,4,9-tetrahydro-pyrano[3,4-b]indol-1-yl]-acetic acid.

### ABSTRACT OF THE DISCLOSURE

The invention is directed to novel pyranoindole derivatives and analogs as well as compositions containing the same and to the use thereof for the treatment, prevention or inhibition of viral infections and associated diseases caused by the Hepatitis C virus.